

Larvicidal Activity of a Neem Tree Extract (Azadirachtin) Against Mosquito Larvae in the Republic of Algeria

Abdelouaheb Alouani^{*}, Nassima Rehim, Nouredine Soltani

Laboratory of Biology Animal Application, Biology Department, Faculty of Sciences, University of Badji Mokhtar, 23000 Annaba, Algeria

Abstract

An insecticide containing azadirachtin, a tree (*Azadirachta indica*) extract, was tested against *Culex pipiens* mosquito larvae and pupae in east of the Republic of Algeria under laboratory conditions. First, after treatment of larval stage, LC_{50} and LC_{90} values for Azadirachtin were 0.35 and 1.28 mg/L in direct effect and 0.3-0.99 mg/l in indirect effect, respectively. Second, after treatment of the pupal stage, the LC_{50} and LC_{90} in direct effect were measured as 0.42 -1.24mg/l and in indirect effect was 0.39mg/l-1.14mg/l respectively. Mosquito adult fecundity were markedly decreased and sterility was increased by the Azadirachtin after treatment of the fourth instar and pupal stage. The treatment also prolonged the duration of the larval stage. The results show that the Azadirachtin is promising as a larvicidal agent against *Culex pipiens*, naturally occurring biopesticide could be an alternative for chemical pesticides.

المخلص

تناول هذا البحث دراسة تأثير مبيد نباتي Azadirachtin المستخلص من شجرة *Azadirachta indica*، على يرقات و عذارى البعوض في شرق جمهورية الجزائر في شروط المختبر، حيث حسبت الجرعة المميتة للنصف (LC_{50}) والجرعة المميتة لتسعين بالمئة (LC_{90}) بالنسبة لبعوض *Culex pipiens* وهي أكثر العوامل نقلا لبعض الأمراض الإلتهابية على الصعيد المحلي، حيث قدرت بعد معاملة الطور الرابع $LC_{50} = 0.35$ ملغ /ل و $LC_{90} = 1.28$ ملغ/ل هذا بالنسبة إلى التأثير المباشر، أما التأثير الغير مباشر فكانت 0.30 ملغ/ل و 0.99 ملغ/ل على التوالي، أما عند معاملة طور العذاراء كانت النتائج بالنسبة للتأثير المباشر $LC_{50} = 0.42$ ملغ /ل و $LC_{90} = 1.24$ ملغ/ل أما التأثير الغير مباشر فكانت 0.39 ملغ/ل و 1.14 ملغ/ل على التوالي. كذلك لوحظ انخفاض كبير في نسبة الخصوبة وزيادة مؤشر العقم للإناث الناتجة عن معاملة الطور الرابع والعذارى للبعوض، وزيادة في مدة الطور. حيث نقول في الأخير أنه يمكن استعمال هذا المبيد كمبيد طبيعي بدل المبيدات الكيميائية.

© 2009 Jordan Journal of Biological Sciences. All rights reserved

Keywords: Mosquito; *Culex pipiens*; Azadirachtin; Insecticide.

1. Introduction

The Meliaceae plant family is known to contain a variety of compounds that show insecticidal, antifeedant, growth-regulating, and development-modifying properties (Nugroho et al., 1999; Greger et al., 2001; D'Ambrosio and Guerriero, 2002; Nakatani et al., 2004). *Melia azedarach* L. and *Azadirachta indica* (Sapindales: Meliaceae), commonly known as Chinaberry or Persian lilac tree, are deciduous trees that are native to northwestern India; and have long been recognized for their insecticidal properties. These trees typically grow in the tropical and subtropical parts of Asia, but nowadays they are also cultivated in other warm regions of the world because of their considerable climatic tolerance. Fruit extracts of *Melia azedarach* and *Azadirachta indica* elicit a variety of effects in insects such as antifeedant, growth retardation, reduced fecundity, moulting disorders, morphogenetic defects, and changes of behavior (Schmidt et al., 1998; Abou Fakhr Hammad et al., 2001; Gajmer et al., 2002; Banchio et al., 2003; Wandscheer et al., 2004). The effects of the compounds extracted from *M. azedarach* on insects

have been reviewed by Ascher et al., (1995) and reported by Saxena et al., (1984), Schmidt et al., (1998), Juan et al., (2000), Carpinella et al., (2003), Senthil Nathan and Saehoon, (2005). Control of mosquito is essential as many species of mosquitoes are vectors of malaria, filariasis, and many arboviral diseases; and they constitute an intolerable biting nuisance (Youdeowei and Service, 1983; Curtis, 1994; Collins and Paskewitz, 1995). Biotechnologists and entomologists agree that mosquito control efficiency should be with selectivity for a specific target organism. New control methodologies aim at reducing mosquito breeding sites and biting activity by using a combination of chemical-biological control methods soothing several advocated biocontrol methods to reduce the population of mosquito and to reduce the man-vector contact (Service, 1983). Recently, there has been a major concern for the promotion of botanicals as environment friendly pesticides, microbial sprays, and insect growth regulators amidst other control measures such as beneficial insects and all necessitate an integration of supervised control (Ascher et al., 1995; Senthil Nathan et al., 2004, 2005b, c, and d). The development of insects' growth regulators (IGR) has received considerable attention for selective control of insect for medical and veterinary importance and has produced mortality due to their high neurotoxic effects (Wandscheer et al., 2004; Senthil Nathan et al.,

^{*} Corresponding author. alouanitoxci@yahoo.fr.

2005a). Although, biological control has an important role to play in modern vector control programs, it lacks the provision of a complete solution by itself. Irrespective of the less harmful and eco-friendly methods, suggested and used in the control programmers, there is still a need to depend upon the chemical control methods in situations of epidemic outbreak and sudden increase of adult mosquitoes. Hence, insecticides are known for their speedy action and effective control during epidemics. Nonetheless, they are preferred as effective control agent to reduce the mosquito population irrespective of their side effects. Recent studies stimulated the investigation of insecticidal properties of plant-derived extracts; and concluded that they are environmentally safe, degradable, and target specific (Senthil Nathan and Kalaivani, 2005). Muthukrishnan and Puspalatha (2001) evaluated the larvicidal activity of extracts from *Calophyllum inophyllum* (Clusiaceae), *Rhinacanthus nasutus* (Acanthaceae), *Solanum suratense* (Solanaceae) and *Samadera indica* (Simaroubaceae), *Myriophyllum spicatum* (Haloragaceae) against *Anopheles stephensi* (Senthil Nathan et al., 2006). Several indigenous plants in India and subtropical parts of Asia, such as *Ocimum basilicum*, *Ocimum santum*, *Azadirachta indica*, *Lantana camera*, *Vitex negundo* and *Cleome viscosa* (Senthil Nathan et al., 2006) were studied for their larvicidal action on the field which collected fourth instar larva of *Culex quinquefasciatus* (Kalyanasundaram and Dos, 1985). Chavan (1984), Zebitz(1984,1986), Schmutterer (1990), Murugan and Jeyabalan (1999) reported that *Leucas aspera*, *O. santum*, *Azadirachta. indica*, *Allium sativum* and *Curcuma longa* had a strong larvicidal, antiemergence, adult repellency and antireproductive activity against *A. stephensi*. In addition, *Pelargonium citrosa* (Jeyabalan et al., 2003), *Dalbergia sissoo* (Ansari et al., 2000a) and *Mentha piperita* (Ansari et al., 2000b) were shown to contain larvicidal and growth inhibitory activity against *A. stephensi*. The present investigation was conducted to study the effect of Azadirachtin, a neem tree *Azadirachta indica* extract, against larvae and pupae of *Culex pipiens* mosquitoes in east of the Republic of Algeria.

2. Materials and Methods

2.1. Mosquito Rearing

Culex pipiens eggs were collected from cellarage tribes (region sidi amar - Annaba) and reared in the ' Laboratory of Biology Animal Application' University of Annaba-Algeria. Larvae were reared in plastic and enamel trays in tap water. They were maintained at 25-27°C, 75-85% relative humidity under 14:10 light and dark photo period cycle. The larvae were fed with fresh food consisting of a mixture of Biscuit Petit Regal-dried yeast (75:25 by weight). Pupae were transferred from the trays to a cup containing tap water and placed in screened cages (20x20x20cm), where the adult emerged. After emergence, female mosquitoes obtained blood meal from caged pigeons while male mosquitoes were fed on a 10% sucrose solution. Then egg-masses were kept to continue next generation.

2.2. Bioassays and Larval Mortality

Bioassays were performed with fourth larvae stages and pupae of *C. pipiens* using concentration from 0.125; 0.250; 0.500; 0.750 and 1mg/l of the Azadirachtin. A minimum of 25 larvae/concentration were used for all the experiments. And these experiment were replicated five times. For mortality studies, 25 larvae each of fourth instar and pupae were introduced in 250 ml plastic beaker containing various concentrations of the Azadirachtin. A control was maintained. The treatments were replicated five times, and each replicate set contained one control. The percentage mortality was calculated by using the formula (1), and corrections for mortality when necessary were done using Abbot's (1925) formula (2)

Percentage of Mortality

$$= \frac{\text{Number of dead larvae}}{\text{Number of larvae introduced}} \times 100 \quad (1)$$

Corrected percentage of mortality

$$= 1 - \frac{n \text{ in T after treatment}}{n \text{ in C after treatment}} \times 100 \quad (2)$$

Where n = number of larvae or nymph, T = treated, C = control.

2.3. Fecundity and Sterility

The fecundity experiments were conducted by taking equal number of male and female *C. pipiens* which had emerged from the control and treated sets of each concentration. They were mated in the cages of (20 x 20 x 20) cm dimension individually to each concentration. Three days after the blood meal, eggs were collected daily from the small plastic bowls containing water kept in ovitraps in the cages. The fecundity was calculated by the number of the eggs laid in the ovitraps divided by the number of female let to mate (The death of the adult in the experiment was also considered) The Sterility Indices experiments were conducted by the formula (3) of Sexina et al.(1993):

$$SI = 100 - \frac{\text{Total number of eggs of females treated} \times \text{percentage of hatch}}{\text{Total number of eggs of control} \times \text{percentage of hatch}} \quad (3)$$

2.4. Total Larval and Pupal Duration

To assay the growth factors of *C. pipiens*, test solution of concentration of Azadirachtin extract (0.125; 0.250; 0.500; 0.750 and 1mg/l) were used. A known number of eggs were made to hatch and the total larval duration (days) was calculated from hatching to pupation period, the pupa was placed in a small container closed with a transparent mesh, so that the adults were kept trapped. The pupal duration (days) was calculated from the pupal molt to the emergence of imago.

2.5. Statistical Analysis

The analysis program Probit (Finney, 1971), the lethal concentrations ($\mu\text{g/ml}$) for 50%, and 90% of the mortality, LC_{50} and LC_{90} , respectively, were at 24h after treatment.

Table 1. Larvicidal activity of Azadirachtin at various concentration applied for 24h to newly ecdysed fourth instars of *Culex pipiens*.

Effects	Concentration (mg/l)	Mortality (%)	LC ₅₀ (mg/l)	95%Confidancelimits (µg/ml)		LC ₉₀ (mg/l)	Regression equation	χ^2
				Lower	Upper			
Direct	Control	9.6						
	0.125	17.39						
	0.250	33.62	0.357	0.307	0.414	1.280	Y=2.20x-0.61	0.615
	0.500	60.17						
	0.750	76.39						
	1	85.24						
Indirect	Control	15.2						
	0.125	19.80						
	0.250	35.52	0.304	0.286	0.335	0.992	Y=2.63x-1.60	3.69
	0.500	65.40						
	0.750	81.12						
	1	93.70						

Table 2. Larvicidal activity of Azadirachtin at various concentration, applied for 24h to newly ecdysed pupae of *Culex pipiens*.

Effects	Concentration (mg/l)	Mortality (%)	LC ₅₀ (mg/l)	95%Confidancelimits (mg/l)		LC ₉₀ (mg/l)	Regression equation	χ^2
				Lower	Upper			
Direct	Control	4.33						
	0.125	7.57						
	0.250	28.78	0.426	0.379	0.479	1.243	Y=2.76x-2.2	1.1625
	0.500	45.45						
	0.750	66.66						
	1	87.87						
Indirect	Control	4						
	0.125	7.93						
	0.250	31.74	0.398	0.355	0.445	1.141	Y=2.82x-2.3	1.485
	0.500	49.20						
	0.750	71.42						
	1	92.05						

The 95% confidence intervals, values, and degrees of freedom of the χ^2 goodness of fit tests, and regression equations, were recorded. Whenever the goodness of χ^2 was found to be significant ($p < 0.05$), a heterogeneity correction factor was used in the calculation of confidence limits. Data from biology, mortality, fecundity deterrence and effective concentration were subjected to analysis of variance (ANOVA of arsine square root transformed percentages).

3. Results

3.1. Insecticidal Activity

Dose-response relationship was determined for Azadirachtin applied for 24h to newly ecdysed fourth instar larvae and pupae. The mortality was scored up to adult formation.

For fourth stage, the highest concentration tested 1 mg/l in direct effect caused 85.24% mortality (Figure1). With probit, the LC₅₀ was calculated as 0.35 mg/l (95%

CI=0.30-0.41mg/l; n=75; Slope=2.83) and the LC₉₀ was 1.28 mg/l, (Table1). For indirect effects, the highest concentration caused 93.70% mortality (Figure2), the LC₅₀ was 0.32 mg/l (95% CI=0.28-0.33 mg/l; n=75; Slope=2.46) and LC₉₀ was calculated as 0.99 mg/l, respectively, (Table1). After treatment, during the pupal stage and in direct effect, the highest concentration caused 87.87% mortality (Figure1), the LC₅₀ was 0.42 mg/l (95% CI=0.37-0.47mg/l; n=75; Slope=2.83) and the LC₉₀ was calculated as 1.24 mg/l, (Table 2). For indirect effects the LC₅₀ was 0.39 mg/l (95% CI=0.35-0.44 mg/l; n=75; Slope=2.46) and LC₉₀ was calculated as 0.99 mg/l respectively, the highest concentration tested caused 92.05% mortality (Table2, Figure2).

3.2. Effect on Fecundity, Sterility and growth

Adults fecundity also was markedly decreased by the Azadirachtin treatment the fourth instar larvae and pupal stages (Figure 3, 4). Adults' sterility indices were markedly increased by the Azadirachtin treatment, (Figure 5, 6). An adverse sub-lethal effect in pupa exposed to Azadirachtin

was evident. In addition to significantly lower survivorship and protracted development, larval duration was reduced markedly. The plant extracts (Azadirachtin) drastically reduced the fecundity of the females, and only few adults survived. The duration of larval instars and the total developmental time were prolonged. (Table3). In the present study, application of Azadirachtin greatly affected the growth of *Culex pipiens*.




Figure1. Dose-response relationship for treatment of Azadirachtin, applied for 24h to newly ecdysed fourth instar larvae and pupae of *Culex pipiens* (effect direct).




Figure 2. Dose-response relationship for treatment of Azadirachtin, applied for 24h to newly ecdysed instar larvae and pupae instars of (*Italiqie*) *Culex pipiens* (effect indirect).

4. Discussion

Azadirachtin, the extract of neem tree, was tested in the present study, and is reported to be eco-friendly and not toxic to vertebrates (Al- Sharook et al., 1991). It is clearly proved that crude or partially-purified plant extracts are less expensive and highly efficacious for the control of mosquitoes rather than the purified compounds or extracts (Jang et al., 2002; Cavalcanti et al., 2004).




Figure 3. Fecundity of *C. pipiens* females after treated the fourth instars with Azadirachtin. (Data following by *** are significantly different from control, $p < 0.001$)




Figure 4. Fecundity of (*Italiqie*) *Culex pipiens* females after treating the pupae stages with Azadirachtin. (Data following by *** are significantly different from control, $p < 0.001$).




Figure5. Sterility Indices of *C. pipiens* females after treating the fourth instars with Azadirachtin. (Data following by *** are significantly different from control, $p < 0.001$).




Figure 6. Sterility Indices of *C. pipiens* females after treating the pupae with Azadirachtin. (Data following by *** are significantly different from control, $p < 0.001$)

Table 3. Effect of Azadirachtin applied to newly ecdysed 4th instar larvae of *C. pipiens* on the duration of development.

Concentration (mg/l)	Duration (days) Mean ± SE	
	4 th larval instar	Pupal stage
0	8.75 ± 2.99	2.61 ± 0.65
0.125	15.24 ± 1.28	3.81 ± 0.84
0.250	16.00 ± 2.82	3.54 ± 0.55
0.500	17.94 ± 1.54	3.42 ± 0.87
0.750	18.45 ± 2.80	3.33 ± 0.28
1	19.75 ± 2.22	3.33 ± 0.46

The effect of these crude plant extract on the biology, reproduction, and adult emergence of the mosquitoes are remarkably greater than those reported for other plant extracts in the literature. For example 50% inhibition of the emergence of the adult mosquitoes was observed by the use of *C. inophyllum*, *S. suratense*, *S. indica* and *Rhinocanthus nasutus* leaf extracts (Muthukrishnan and Puspapalatha, 2001). Similarly 88% of the adult mortality was observed by the use of *P. citrosa* leaf extracts at 2% concentration (Jeyabalan et al., 2003). The Meliaceae plant family is used as growth regulator against many insect pests (Saxena et al., 1984; Jacobson, 1987; Schmutterer,

1990; Hammad et al., 2001; Gajmer et al., 2002; Banchio et al., 2003; Wandscheer et al., 2004). The growth regulatory effect is the most important physiological effect of *M. azedarach* on insects. It is because of this property that family Meliaceae has emerged as a potent source of insecticides. Exposure of *A. stephensi* larvae to sub-lethal doses of neem leaves extract in the laboratory prolonged larval development, reduced pupal weight and oviposition (Murugan et al., 1996; Su and Mulla, 1999). In the field, delayed phenology of surviving larvae and reduced pupal weight are common occurrence after treatment with neem (Zebitz, 1984, 1986). The direct and indirect contribution of such effects to treatment efficacy through reduced larval feeding and fitness need to be properly understood in order to improve the use of *M. azedarach* for management of *A. stephensi*. The results of this study indicate the plant-based compounds such as Azadirachtin (compounds present in the Meliaceae plant family seed) may be an effective alternative to conventional synthetic insecticides for the control of *Culex pipiens*. Undoubtedly, plant derived toxicants are valuable sources of potential insecticides. These and other naturally occurring insecticides may play a more prominent role in mosquito control programs in the future (Mordue and Blackwell, 1993). The results of this study will contribute to a great reduction in the application of synthetic insecticides, which in turn will increase the opportunity for natural control of various medically important pests by botanical pesticides. Since these are often active against a limited number of species including specific target insects, less expensive, easily biodegradable to non-toxic products, and potentially suitable for use in mosquito control programme (Alkofahi et al., 1989), they could lead to development of new classes of possible safer insect control agents. Plant allelochemicals may be quite useful in increasing the efficacy of biological control agents because plants produce a large variety of compounds that increase their resistance to insect attack (Berenbaum, 1988; Murugan et al., 1996; Senthil Nathan et al., 2005a). The intensive use of pesticides produces side effects on many beneficial insects and also poses both acute and chronic effects to the milieu (Abudulai et al., 2001). Recently, bio-pesticides with plant origins are given for use against several insect species especially disease-transmitted vectors, based on the fact that compounds of plant origin are safer in usage, without phytotoxic properties; also leave no scum in the environment (Schmutterer, 1990; Senthil Nathan et al., 2004, 2005a, d). Large alterations in the fecundity and sterility of insects exposed to neem have been extensively reported, such as those in the fly, *Ceratitis capitata* (Steffens and Schmutterer 1982); banana root borer, *Cosmopolites sordidus* (Germar) (Musabyimana et al., 2001); and mosquitoes, *A. stephensi* and *A. culicifacies* (Dhar et al., 1996). The work published by Khan et al., (2007) microscopically demonstrated that the decrease in fecundity of *Bactocera cucurbitae* and *Bactocera dorsalis* exposed to neem compound was due to the block of ovarian development. Likewise, mixing of a commercial formulation of neem in the adult diet caused reduction in the fecundity of *C. capitata* by interfering with oogenesis (Di Ilio et al., 1999). The block in the ovarian activity of *C. capitata*, resulting from neem compound, was verified by histological observation (Di Ilio et al., 1999). Results

from the study of Lucantoni et al., (2006) clearly indicated that the neem treated female mosquito, *A. stephensi*, displayed a delay in oocyte development in the vitellogenesis. As discussed by Weathersbee III and Tang (2002), the disruption of reproductive capability could lead to substantial population decline over time. Furthermore, Dhar et al., (1996) revealed that exposure to neem extract suppressed rather than inhibited oviposition in mosquitoes. The present study clearly proved the efficacy of Azadirachtin on larvae, pupae, and adult of *Culex pipiens*. Further studies such as mode of action and synergism with the biocides under field condition are needed.

Acknowledgements

The authors are grateful to Professor G. Smagghe (Laboratory of Agrozoology, Department of Crop Protection, Faculty of Bioscience Engineering, Ghent, University B-9000 Ghent, Belgium). This research was supported by the Algerian Agency for Research Development in Health (Project 01/07/08/00/02).

References

- Abbot WS. 1925. A method of computing the effectiveness of an insecticide. *J. Econ. Ent.* 18, 265–267.
- Abou Fakhr Hammad M, Zournajian H, Talhouk S. 2001. Efficacy of extracts of *Melia azedarach* L. callus, leaves and fruits against adults of the sweet potato whitefly *Bemisia tabaci* (Homoptera: Aleyrodidae). *J. Appl. Entomol.* 125, 483–488.
- Abudulai M, Shepard BM, Mitchell PL. 2001. Parasitism and predation on eggs of *Leptoglossus phyllopus* (L.) (Hemiptera: Coreidae) in cowpea: impact of endosulfan sprays. *J. Agric. Urban Entomol.* 18, 105–115.
- Alkofahi A, Rupprecht JK, Anderson JE, Mclaughlin JL, Mikolajczak KL, Scott BA. 1989. Search for new pesticides from higher plants. In: Arnason, JT., Philogene, BJR., Morand, P. (Eds.), *Insecticides of Plant Origin*. American Chemical Society, Washington, DC, pp. 25–43.
- Al-Sharook Z, Balan K, Jiang Y, Rembold H, 1991. Insect growth inhibitors from two tropical meliaceae. Effect of crude seed extracts on mosquito larvae. *J. Appl. Ent.* 111, 425–430.
- Ansari MA, Razdan RK, Tandon M, Vasudevan P. 2000a. Larvicidal and repellent actions of *Dalbergia sissoo* Roxb. (F. Leguminosae) oil against mosquitoes. *Biores. Technol.* 73, 207–211.
- Ansari, MA, Vasudevan, P, Tandon, M, Razdan, RK. 2000b. Larvicidal and mosquito repellent action of peppermint (*Mentha piperita*) oil. *Biores. Technol.* 71, 267–271.
- Ascher KRS, Schmutterer H, Zebitz CPW, Naqvi SNH. 1995. The Persian lilac or Chinaberry tree: *Melia azedarach* L. In: Schmutterer, H. (Ed.), *The Neem Tree: Source of Unique Natural Products for Integrated Pest Management, Medicine, Industry and Other Purposes*. VCH, Weinheim, Germany, pp. 605–642.
- Banchio E, Valladares G, Defago M, Palacios S, Carpinella, C. 2003. Effects of *Melia azedarach* (Meliaceae) fruit extracts on the leafminer *Liriomyza huidobrensis* (Diptera: Agromyzidae): assessment in laboratory and field experiments. *Ann. Appl. Biol.* 143, 187–193.
- Berenbaum MR. 1988. Allelochemicals in insect–microbe–plant interactions: agents provocateurs in the coevolutionary arms race. In: Barbosa, P, Lotourneau, DK. (Eds.), *Novel Aspects of Insect–Plant Interactions*. Wiley, New York, pp. 97–123.

- Carpinella MC, Defago MT, Valladares G, Palacios SM. 2003. Antifeedant and insecticide properties of a limonoid from *Melia azedarach* (Meliaceae) with potential use for pest management. *Agric. Food Chem.* 15 (51), 369–674.
- Cavalcanti ESB, de Moraes SM, Ashley ALM, William PSE. 2004. Larvicidal activity of essential oils from Brazilian plants against *Aedes aegypti* L. *Memo'rias do Instituto Oswaldo Cruz* 99, 541–544.
- Chavan FR. 1984. Chemistry of alkanes separated from leaves of *Azadirachta indica* and their larvicidal/ insecticidal activity against mosquitoes. In: *Proceedings of 2nd International Neem Conference, Rauschholzhausen*, pp.59-66.
- Collins FH, Paskewitz SM. 1995. Malaria: current and future prospects for control. *Ann. Rev. Entomol.* 40, 195–219.
- Curtis CF. 1994. Should DDT continue to be recommended for malaria vector control?. *Med. Vet. Entomol.* 8 107–112.
- D'Ambrosio M, Guerriero A. 2002. Degraded limonoids from *Melia azedarach* and biogenetic implications. *Phytochemistry* 60, 419– 424.
- Dhar R, Dawar H, Garg S, Basir SF, Talwar G.P .1996. Effect of volatiles from neem and other natural products on gonotrophic cycle and oviposition of *Anopheles stephensi* and *An. Culicifacies* (Diptera: Culicidae). *J Med Entomol* 33:195–201.
- Di Ilio V, Cristofaro M, Marchini D, Nobili P, Dallai R .1999. Effects of a neem compound on the fecundity and longevity of *Ceratitis capitata* (Diptera: Tephritidae). *J Econ Entomol* 92:76–82
- Finney DJ. 1971. *Probit Analysis*, third ed. Cambridge University Press, London, UK, p. 38.
- Gajmer T, Singh R, Saini RK, Kalidhar S.B. 2002. Effect of methanolic extracts of neem (*Azadirachta indica* A. Juss) and bakain (*Melia azedarach* L.) seeds on oviposition and egg hatching of *Earias vittella* (Fab.) (Lepidoptera: Noctuidae). *J. Appl. Entomol.* 126, 238–243.
- Greger H, Pacher T, Brem B, Bacher M, Hofer O. 2001. Flavaglines and other compounds from Fijian *Aglaia* species. *Phytochemistry* 57, 57–64. 1322 S.S.
- Jacobson M. 1987. Neem research and cultivation in Western hemisphere. In: Schmutterer, H., Ascher, KRS. (Eds.), *Natural Pesticide from the Neem Tree and Other Tropical Plants*. *Proceedings of the 3rd Neem Conference, Nairobi, Kenya*, pp. 33–44.
- Jang YS, Kim MK, Ahn YJ, Lee HS.2002. Larvicidal activity of Brazilian plants against *Aedes. aegypti* and *Culex pipiens pallens* (Diptera: Culicidae). *Agric. Chem. Biotechnol.* 44, 23–26.
- Jeyabalan D, Arul, N, Thangamathi P. 2003. Studies on effects of *Pelargonium citrosa* leaf extracts on malarial vector, *Anopheles stephensi* Liston. *Biores. Tech.* 89 (2), 185 189.
- Juan A, Sans A, Riba M. 2000. Antifeedant activity of fruit and seed extracts of *Melia azedarach* and *Azadirachta indica* on larvae of *Sesamia nonagrioides*. *Phytoparasitica* 28, 311–319.
- Kalyanasundaram M, Dos PK. 1985. Larvicidal and synergistic activity of plant extracts for mosquito control. *Ind. J. Med. Res.* 82, 1–19.
- Khan M, Hossain MA, Islam M.S .2007. Effects of neem leaf dust and a commercial formulation of a neem compound on the longevity, fecundity and ovarian development of the melon fly, *Bactocera cucurbitae* (Coquillett) and the oriental fruit fly, *Bactrocera dorsalis* (Hendel) (Diptera: Tephritidae). *Pak J Biol Sci* 10:3656– 3661.
- Lucantoni L, Giusti F, Cristofaro M, Pasqualini L, Esposito F, Lupetti P, Habluetzel A .2006. Effects of a neem extract on blood feeding, oviposition and oocyte ultrastructure in *Anopheles stephensi* Liston (Diptera: Culicidae). *Tissue Cell* 38:361–371.
- Mordue (Luntz) AJ, Blackwell A. 1993. Azadirachtin an update. *J. Insect Physiol.* 39, 903–924.
- Murugan K, Jeyabalan D. 1999. Mosquitocidal effect of certain plants extracts on *Anophels stephensi*. *Curr. Sci.* 76, 631–633.
- Murugan K, Babu R, Jeyabalan D, Senthil Kumar N, Sivaramakrishnan S. 1996. Antipupational effect of neem oil and neem seed kernel extract against mosquito larvae of *Anopheles stephensi* (Liston). *J. Ent. Res.* 20, 137–139.
- Musabyimana T, Saxena RC, Kairu EW, Ogo CPKO, Khan ZR .2001. Effects on neem seed derivatives on behavioral and physiological responses of the *Cosmopolites sordidus* (Coleoptera: Curculionidae). *J Econ Entomol* 94:449–454.
- Muthukrishnan J, Puspalatha E. 2001. Effects of plant extracts on fecundity and fertility of mosquitoes. *J. Appl. Entomol.* 125, 31–35.
- Nakatani M, Abdelgaleil SAM, Saad MMG, Huang RC, Doe N, Iwagawa T. 2004. Phragmalin limonoids from *Chukrasia tabularis*. *Phytochemistry* 65, 2833–2841.
- Nugroho BW, Edrada RA, Wray V, Witte L, Bringmann G, Gehling M, Proksch P. 1999. An insecticidal rocaglamide derivatives and related compounds from *Aglaia odorata* (Meliaceae). *Phytochemistry* 51, 367–376.
- Saxena RC, Epino PB, Cheng-Wen T, Puma BC. 1984. Neem, chinaberry and custard apple: antifeedant and insecticidal effects of seed oils on leafhopper and planthopper pests of rice. In: *Proceedings of 2nd International Neem Conference, Rauschholzhausen, Germany*, pp. 403–412.
- Saxena RC, Harshan V, Saxena A, Sukumaran P.1993. Larvicidal and chemosterilant activity of *Annona squamosa* alkaloids against *Anophel stephensi*. *J. Amer. Mosq. Control. Assoc.* 9: 84-87
- Schmidt GH, Rembold H, Ahmed AAI, Breuer AM. 1998. Effect of *Melia azedarach* fruit extract on juvenile hormone titer and protein content in the hemolymph of two species of noctuid lepidopteran larvae (Insecta: Lepidoptera: Noctuidae). *Phytoparasitica* 26, 283–291.
- Schmutterer H. 1990. Properties and potential of natural pesticides from the neem tree, *Azadirachta indica*. *Ann. Rev. Ent.* 35, 271–297.
- Service MW. 1983. Management of vector. In: Youdeowei, A., Service, N. (Eds.), *Pest and Vector Management in the Tropics*. Longman group Ltd., England, p. 7, 20.
- Senthil Nathan S. Kalaivani K. 2005. Efficacy of nucleopolyhedrovirus (NPV) and azadirachtin on *Spodoptera litura Fabricius* (Lepidoptera: Noctuidae). *Biol. Control.* 34, 93–98.
- Senthil Nathan S, Saehoon K. 2005. Effects of *Melia azedarach* L. extract on the teak defoliator *Hyblaea puera Cramer* (Lepidoptera: Hyblaeidae). *Crop Prot.*, in press.
- Senthil Nathan S, Chung PG, Murugan K. 2004. Effect of botanicals and bacterial toxin on the gut enzyme of *Cnaphalocrocis medinalis*. *Phytoparasitica.* 32, 433–443.
- Senthil Nathan S, Kalaivani K, Murugan, K, Chung P.G. 2005a. The toxicity and physiological effect of neem limonoids on *Cnaphalocrocis medinalis* (Guene'e), the rice leaf folder. *Pest. Biochem. Physiol.* 81, 113–122.
- Senthil Nathan S, Chung P.G, Murugan K. 2005b. Effect of biopesticides applied separately or together on nutritional indices of the rice leaf folder *Cnaphalocrocis medinalis* (Guene'e) (Lepidoptera: Pyralidae). *Phytoparasitica* 33, 187–195.

- Senthil Nathan S, Kalaivani K, Murugan K, Chung PG. 2005c. Efficacy of neem limonoids on *Cnaphalocrocis medinalis* (Guene'e) (Lepidoptera: Pyralidae) the rice leaffolder. *Crop Prot.* 24, 760–763.
- Senthil Nathan S, Kalaivani K, Chung P.G. 2005d. The effects of Azadirachtin and Nucleopolyhedrovirus (NPV) on midgut enzymatic profile of *Spodoptera litura* Fab. (Lepidoptera: Noctuidae). *Pest. Biochem. Physiol.*, in press.
- Senthil Nathan S, Savitha G, George DK, Narmadha A, Suganya L, Chung PG. 2006. Efficacy of *Melia azadirach* L. Extract on the malarial vector *Anopheles Stephensi* Liston (Diptera: Culicidae). *Bioresource Technology*. 97, 1316-1323.
- Steffens RJ, Schmutterer H. 1982. The effect of a crude methanolic neem (*Azadirachta indica*) seed kernel extract on metamorphosis and quality of adults of the Mediterranean fruit fly, *Ceratitis capitata* Wied. (Diptera: Tephritidae). *Z Angew Entomol* 94:98–103.
- Su T, Mulla M.R. 1999. Oviposition bioassay responses of *Culex tarsalis* and *Culex quinquefasciatus* to neem products containing azadirachtin. *Entomol. Exp. Appl.* 91, 337–345.
- Wandscheer CB, Duque JE, da Silva MAN, Fukuyama Y, Wohlke JL, Adelman J, Fontana JD. 2004. Larvicidal action of ethanolic extracts from fruit endocarps of *Melia azedarach* and *Azadirachta indica* against the dengue mosquito *Aedes aegypti*. *Toxicon* 44, 829–835.
- Weathersbee III A.A, Tang Y.Q. 2002. Effect of neem extract on feeding, growth, survival, and reproduction of *Diaprepes abbreviatus* (Coleoptera: Curculionidae). *J Econ Entomol* 95:661–667.
- Youdeowei T, Service M.W. 1983. *Pest Control and Management*. Longman Singapore Publishers Ltd., Singapore, p. 17.
- Zebitz CPW. 1984. Effects of some crude and azadirachtin enriched neem *Azadirachta indica* seed kernel extracts on larvae of *Aedes aegypti*. *Entomol. Exp. Appl.* 35, 11–14.
- Zebitz CPW. 1986. Effects of three neem seed kernel extracts and azadirachtin on larvae of different mosquito species. *J. Appl. Entomol.* 102, 455–463.

