

The Role of Silymarin in the Protection of Mice Liver Damage Against Microcystin-LR Toxicity

Wahsha M. , Al-Jassabi S.*

Department of Biological Sciences, Yarmouk University, Jordan

Abstract

The presence of cyanobacterial toxins in drinking and recreational waters represents a potential risk to public health. Microcystin-LR (MC-LR) is a potent cyclic heptapeptide hepatotoxin produced by the blue-green algae *Microcystis aeruginosa* (*M. aeruginosa*). Chemoprotectant studies suggest that membrane-active antioxidants may offer a protection against microcystin toxicity. The aim of this study is to investigate the potential benefits of dietary supplementation of silymarin as antioxidant on microcystin toxicity in mouse livers. A group of Balb/c mice was pre-treated for ten days with silymarin (extracted from milk thistle seeds collected from local areas of north Jordan) (400 mg of silymarin /Kg mouse body weight given orally once a day for 10 days), before an intraperitoneal injection (i.p) with 200 µg toxin/kg mouse body weight of MC-LR (according to LD₅₀ value). Pre-treatment of mice with a single dose of silymarin, aflavonolignane (*Silybum marianum* L. Geartin) completely abolished the lethal effects and significantly decreased the levels of serum enzymes, alanine aminotransferase (ALT), lactate dehydrogenase (LDH) and gamma glutamyl transferase (GGT), inhibition of Protein phosphatase (PP1), level of methylglyoxal (MG) and lipid peroxidation (LPO) as MDA amount induced by MC-LR. Therefore, silymarin supplied as dietary supplement may have protective effects against chronic exposure to MC-LR.

المخلص

يشكل وجود سموم الطحالب الزرقاء الخضراء (السيانوبكتريا) في مياه الشرب والري مشكلة صحية خطيرة للغاية. يمتاز هذا النوع من السموم بأنه ذا سمية عالية الكفاءة وخاصة لنسيج الكبد حيث وجد أنه يسبب تلف نسيج الكبد بسرعة فائقة إضافة إلى ما يسببه من أورام سرطانية. وهذا النوع من السموم الطحلبية هو عبارة عن بيتيد سباعي حلقي يقوم بتكوينه نوع من الطحالب الزرقاء -الخضراء والمسماة بالميكروسستس اريجينوزا. وقد اثبتت الدراسات الخاصة باستخدام الغذائية التي تساعد في حماية الانسجة الحية بان مضادات الاكسدة قد تساعد في الحماية من هذا النوع من السموم. ان هدف هذه الدراسة هو البحث عن امكانية استخدام مضادات الاكسدة الغذائية كالسليمارين (المستخلص من نبات حليب مريم الموجود في المناطق الشمالية من المملكة) في حماية كبد الفأر من التلف الذي تسببه له هذه السموم. لقد تم اخذ مجموعة من الفئران من نوع Balb/c وتم اطعمها بمستخلص السليمارين بمقدار 400 ملغم لكل كغم من جسم الفأر ولمدة عشرة ايام (مرة واحدة يوميا ولمرة واحدة) ومن ثم حقنها في اليوم الحادي عشر بسموم المايكروسستين تحت الجلد بجرعة مقدارها 200 مايكروغرام لكل كغم من وزن الفأر (وذلك نسبة الى قيمة LD50) التي تم قياسها. ومن ثم تم قياس كفاءة السليمارين عن طريق قياس كمية كل من الانزيمات الالانين ترانسفيريز، اللاكتيت ديهيدروجينيز والكلوتاميل ترانسفيريز في مصل الدم. وقياس كل من البروتين فوسفاتيز والمثيل كلايوكزاليز، ومقدار اكسدة الليبيدات (قياس كمية المالون ثنائي الالدهيد في مستخلص نسيج الكبد. وقد وجد في هذه الدراسة ان السليمارين يتمتع بكفاءة عالية كمضاد للاكسدة التي يسببها هذا النوع من السموم وبذلك يوفر الحماية المطلوبة لكبد الفأر من التلف.

© 2009 Jordan Journal of Biological Sciences. All rights reserved

Keywords: Chemoprotection; Toxicity; Microcystin; Silymarin.

1. Introduction

The rising eutrophication by human activities of fresh waters, including drinking water reservoirs, has increased the occurrence and intensity of cyanobacterial blooms (Ordorika *et al.* 2004; Antoniou *et al.*, 2008). Microcystins (MCs), specific hepatotoxins produced by numerous cyanobacterial species (primarily *M. aeruginosa*) in eutrophic surface waters have risen, increasing worldwide concern (Andrinolo *et al.*, 2008; Billam *et al.*, 2008). MCs are characterized as monocyclic heptapeptide with over 70 different MCs isoforms identified (Spoof and Meriluoto,

2002). The general structure of microcystin is cyclo (D-Ala-L-X-D-MeAsp-L-Z-Adda-D-Glu-Mdha) where X and Z represent. The two variable amino acids and Adda is 3-amino-9-methoxy-2, 6, 8-trimethyl-10-phenyl-deca-4, 6-dienoic acid (Ordorika *et al.*, 2004; Oriola and Lawton, 2005).

Microcystin-LR (MC-LR) is the most widely distributed and studied MCs variant (Gupta and Guha, 2006). It has the amino acids leucine and arginine at positions 2 and 4, respectively (Spoof and Meriluoto, 2002; Antoniou *et al.*, 2008). MC-LR has been found to be a potent inhibitor of protein phosphatase type 1 and type 2A (Andrinolo *et al.*, 2008), resulting in the disturbance of many important cellular processes (Oriola and Lawton, 2005). Exposure to MC-LR has been shown

* Corresponding author. sjassabio@hotmail.com.

to cause oxidative stress in various organisms (Leao *et al.*, 2008). Formation of reactive oxygen species (ROS) and oxidative stress is associated with the development of many pathological states (Gupta and Guha, 2006; Billam *et al.*, 2008). Oxidative stress may occur either due to the decrease of cellular antioxidant level, or due to the overproduction of ROS (Jayaraj *et al.*, 2006). Exposure to MC-LR has been linked with increase of ROS production not only in domestic and wildlife animals, but also in human (Weng *et al.*, 2007).

Recently, interest has increased considerably in finding naturally occurring antioxidants to replace synthetic antioxidants, which are being restricted due to their carcinogenicity (Looa *et al.*, 2007). Silymarin was shown to be an antihepatotoxic agent against MC-LR toxicity in cultured rat hepatocytes (Mereish and Solowe, 1990). Mereish *et al.*, (1991) extended their studies to *in vivo* and tested the use of silymarin against MC-LR toxicity in mice and rats, found that this compound acted potentially against it. Rao *et al.*, (2004) concluded that silymarin could offer great protection against MC-LR toxicity.

The goal of this study is to investigate the role of naturally- extracted silymarin (one of the active polyphenol antioxidants in milk thistle seeds) pre-treatment in protection against MC-LR (extracted from King Talal Reservoir, Jordan)-induced liver injury in Balb/c mice.

2. Materials and Methods

2.1. Chemicals

All chemicals used in this study were of analytical grade and were purchased from Sigma Chemical Co. USA.

2.2. Samples

Samples of *M. aeruginosa* cells were collected according to (Al-Jassabi and Khalil, 2006) from selected sites of KTR in Jordan during blooming season (July, August and September, 2007). Microcystin was extracted from the lyophilized cells of *M. aeruginosa* according to Lawton *et al.*, (1994), and the LD₅₀ of the toxin extract was determined according to Fawell *et al.*, (1999), in which LD₅₀ is determined as the value between the lowest toxin dose at which mice die and the highest toxin dose at which mice live after administration of a toxin.

2.3. Silymarin Extraction

Milk thistle seeds were collected from specific areas in Jordan (Ajlun & Irbid) during July and August 2007. The silymarin was extracted according to the method reported by Duan *et al.*, (2004).

2.4. Animal Treatment and Sample Collection

Male Balb/c mice 6-7 weeks old (average body weight 30 g) were used in this study. Mice were obtained from animal house/ Yarmouk University and were maintained on standard laboratory diet and tap water throughout the experiments. Five animals were maintained in each cage (stainless) under a 12-12h light-dark cycle and room temperatures of 23-26°C.

Sixty mice were used in this study, and they were assigned into 4 groups (10 mice each) as follows: Group 1 was the control group (C), without supplementation of

silymarin or treatment with toxin; Group 2 was the toxin control group (with 2 sub groups, 10 mice each) (T₆ and T₁₂), treated with toxin only, intraperitoneal injection (i.p) with 200 µg toxin / kg mouse body weight. 10 mice were killed after 6 h and 10 mice were killed after 12 h; Group 3 was the silymarin control group (SC), supplemented orally with 400 mg silymarin /kg mouse body weight daily for 10 days according to Lakshmana *et al.*, (2004), then they were killed; Group 4 (silymarin and toxin) with two sub group (10 mice each) supplemented orally with 400 mg silymarin / kg mouse body weight daily for 10 days, then were injected i.p. with 200 µg toxin/kg mouse body weight. 10 mice were killed after 6 h (ST₆) and 10 mice were killed after 12 h (ST₁₂). Blood was collected immediately after sacrifice of mice, and serum was isolated and stored at -20°C for the biochemical tests. Livers were removed, immediately after death, perfused with normal saline containing heparin, weighed, and homogenized with phosphate buffer saline (pH 7.2) using Ultra Turax homogenizer, centrifuged at 3000g for 30min. The supernatant was removed and stored at -20°C.

2.5. Alanine Aminotransferase (ALT) Assay

Determination of ALT activity in the serum sample was measured according to the procedure recommended by Gehringer *et al.*, (2003).

2.6. Cytotoxicity Assay

Serum lactate dehydrogenase (LDH) activity was measured according to Doyle and Griffiths, (1998).

2.7. Gamma Glutamyl Transferase Assay (GGT):

Determination of GGT activity in the serum sample was measured based on Szasz (1974) method .

2.8. Protein Phosphatase (PPI) Assay

Depending on the procedure described by Yuan *et al.*, (2006), PPI activity in liver homogenate was assayed by measuring the rate of formation of the yellow color of p-nitrophenol (p-NP) produced by hydrolysis of p-nitrophenylphosphate (p-NPP) in an alkaline solution spectrophotometrically.

2.9. Lipid Peroxidation (LPO) Assay

The lipid peroxidation level of the hepatocyte was measured according to the method described by Hosseinzadeh *et al.*, (2007).

2.10. Methylglyoxal Assay (MG)

The glyoxalase system breaks down the toxic methylglyoxal, which is formed as a by-product of the triosephosphate isomerase reaction in glycolysis. Methylglyoxal was determined from liver homogenate according to the method described by Ratliff *et al.*, (1996).

2.11. Statistical Analysis

All results were expressed as the mean ± S.E.M from ten mice per group. One way analysis of variance (ANOVA) followed by a Tukey test was used to determine the significance of the differences between the groups. Statistical significance was declared when P value was equal to or less than 0.05. The statistical analysis was performed using the Sigma Stat Statistical Software version 3.5.

3. Results

In this study, MC-LR treated mice showed increased liver body mass index ratio due to massive intrahepatic hemorrhage and pooling of blood in the liver, as shown in Table 1. The livers of mice which received silymarin were within the normal value, but this value increased slightly after the exposure to the toxin.

Treatment with MC-LR resulted in a significant increase in levels of ALT activity compared to the saline-treated group (Table 1). The elevated levels of serum ALT were significantly reduced in mice received toxin and silymarin supplementation. An increased ALT value was revealed in sera by folds of 1.5 in ST₆. A further increase was shown as 4.2, folds for ST₁₂ compared to controls (P<0.05).

As shown in Table 1, MC-LR alone produced after 6 h of exposure nearly three-fold and five-fold after 12 h increases in serum LDH level compared to control mice, indicating severe liver injury. The mice receiving toxin and silymarin supplementation revealed an increased LDH value in sera by folds of 2 in ST₆. A further increase was shown as 2.5 folds for ST₁₂ compared to controls (P<0.05).

Mice group, which received silymarin only, show levels of GGT which are within the normal value (39±4U/mg). Furthermore MC-LR administration (200mg toxin/kg mouse body weight) of 6 and 12 h increased serum values of GGT about 3.8 and 5.8 fold, respectively, compared with control mice (P<0.05), while the increase was dramatically diminished by silymarin pre-treatment, and it could significantly inhibit the increase of GGT induced by MC-LR as shown in Table 1.

Results of spectrophotometric measurements of protein phosphatase activity of liver homogenates for all groups are presented in Table 1. PP1 activity was significantly inhibited in group 2; 0.46% inhibition has occurred for those of 6 h and 72% for those of 12 h when compared with control. Supplementations with silymarin caused a partial protection of PP1 activity against the action of MC-LR, of almost three-fold compared with the levels in toxin control group.

Control mice exhibited normal levels of lipid peroxidation (LPO) measured as amount of MDA it was 0.067µM in hepatocytes homogenate. There was a dramatic increase in MDA level in liver homogenate from toxin-treated groups by 16 fold in case of T₆ and 34 fold in case of T₁₂ when compared with those of control mice (P<0.05), as reflected in the elevation of TBA values, which appear time-dependent (Table 1). The mice receiving toxin and silymarin supplementation revealed an increased TBA value by 3 folds in ST₆ and further increase shown as 10, fold for ST₁₂ in compared to controls.

The MG concentration in the liver homogenates was determined at 6 and 12h after i.p injection of MC-LR and results were compared with controls as shown in Table 1. It was clear that the effect of MC-LR on MG was time dependent; for mice exposed only to the toxin; results were found to be higher at 12 h compared to those at 6 h. However, pre-treated silymarin group showed a significant protection against accumulation of MG in the hepatocytes.

4. Discussion

M. aeruginosa dominates the cyanobacterial communities of KTR during the warmer season (specifically, from June to October, 2007) (AlJassabi, 2004). Our microscopic investigations ensured that the major cyanobacterial species found in the collected samples were *M. aeruginosa* in agreement with our previous study (Al-Jassabi and Khalil, 2006). The phosphatase inhibitory activity proved the bioactivity of the toxin while the spectrophotometric analysis proved that the extracted toxin was MC-LR (Spoof and Meriluoto, 2002 & Oriola and Lawton, 2005).

The present investigation examined the hepatotoxic effects of MC-LR isolated from *M. aeruginosa* of KTR, in Balb/c mice after intraperitoneal route of exposure to the toxin. Besides we investigated the potential hepatoprotective efficacy of silymarin as naturally isolated antioxidant against MC-LR effects. The LD₅₀ concentration of MC-LR by i.p route was determined as 200 µg toxin/kg mouse body weight by a modified Fawell's up-and-down method (Fawell *et al.*, 1999) which is at variation with some earlier reports (Ding *et al.*, 2006 and Lombardo *et al.*, 2006). A wide range of LD₅₀ values have been reported for MC-LR by administration in mice.

In agreement with previous studies (Dawson, 1998; Weng *et al.*, 2007; Xu *et al.*, 2007) the results in toxin group indicate that severe liver damage accompanied by marked change in colour and weight can occur by i.p injection of LD₅₀ dose. Hepatocellular damage was first noticed by the increase in total liver size, due to intrahepatic haemorrhage and accumulation of fluids caused by the action of MC-LR (Dufour and Clavien, 2005).

Exposure to MC-LR causes a disturbance of cellular iron homeostasis as a result of ferritin inhibition (Chen *et al.*, 2005). Mackintosh *et al.*, (1990) reported that Protein phosphatases are inhibited with high affinity by MC-LR. Thus, MC-LR completely blocks access to the active centre of the enzyme using Fe⁺³ as a catalyst (Lohse *et al.*, 1995), thereby causing hyperphosphorylation of the cell and a massive disruption of a number of important cellular mechanisms (Yang *et al.*, 1997). Silymarin in the mice pre-treated group could partially hinder the inhibition of PP1 binding to Fe⁺³ (Kanaze *et al.*, 2005).

One important consequence of excessive free radical production after exposure to MC-LR is the toxin ability to attack many organic molecules, including polyunsaturated fatty acids in the cell membrane (Denisov *et al.*, 2003 and Pinho *et al.*, 2005), leading to lipid peroxidation. Several studies reported that ROS can initiate lipid peroxidation through the action of hydroxyl radicals (Joshi *et al.*, 2005). Our analysis of hepatocytes for lipid peroxidation showed that silymarin could bring a decrease in the formation of lipid peroxidation through their ability to scavenge the hydroxyl radicals. The activity of silymarin as scavenger of free radicals and ROS has been described by others (Mereish *et al.*, 1991; Rao *et al.*, 2004)

Methylglyoxal is considered to be toxic for mammals (Kalapos, 2008). The cytotoxicity associated with the accumulation of MG after exposure to MC-LR is due to the inhibition of glyoxalase I and the depletion of the action of the antioxidant defense system found in the liver

Table 1. Summary of results of the effect of silymarin supplementation on mice receiving single lethal dose of MC-LR

	C	T ₆	T ₁₂	SC	ST ₆	ST ₁₂
Livers weight (g)	1.47 ± 0.02	2 ± 0.057	2.74 ± 0.06	1.45 ± 0.02	1.55 ± 0.05	1.87 ± 0.09
ALT (U/L)	574.6±6.95	1649±43	2461±51	533±5	884±6	1202±39
LDH (U/mg)	987 ± 207	3352± 123	4868 ±46	885 ± 2	2057 ± 67	2471 ± 112
GGT (U/mg)	39 ± 4	132 ± 5	226 ± 7	32 ±4	95 ± 3	163 ±3
PP1 (U/mg)	0.583 ± 0.01	.316± 0.01	0.165±0.002	0.562±0.02	0.414±0.021	0.31±0.007
LPO (µM)	0.067 ± 0.01	1.3 ± 0.04	2.292 ± 0.04	0.013± 0.001	0.212 ± 0.002	0.705 ±0.027
MG (µM)	1.61±0.075	29.03±0.39	41.94±0.797	1.58±0.0411	16.13±0.345	16.94±0.334

(Aljassabi, 2004). The increase in MG levels in the liver caused significant generation of free radicals which might further strengthen the damage and affect the hepatocyte function (Kalapos, 2008). This was noticed in the toxin group which received the toxin only as shown in the results. Silymarin pre-treated group showed a significant decrease in the accumulation of MG.

The antagonistic effect of silymarin could be similar to that of dithioerythritol by stabilizing protein-thiol, which may be important to the structure of liver cell (Mereish et al., 1991). Previous studies and the present investigation show that among the various chemoprotectants that have been screened till date, silymarin is the most effective in preventing the MC-LR-induced lethality in mice.

References

- Al-Jassabi S. 2004. Effect of microcystin from Jordan on ion regulation and antioxidant system in the hepatocytes of mice. *J. Biol. Sci.* 4(4): 547-552.
- Al-Jassabi S. 2004. Effect of methylglyoxal on antioxidant enzymes of the liver and spleen of the mice. *J. Biol. Sci.* 4(5):605-608.
- Al-Jassabi S. and Khalil A. M. 2006. Initial report on identification and toxicity of *Microcystis* in King Talal reservoir, Jordan. *Lakes & Reserv.: Res. & Manag.* 11: 125-9.
- Antoniou MG Shoemaker JA Armah CA and Dionysiou DD. 2008. LC/MS/MS structure elucidation of reaction intermediates formed during the TiO₂ photocatalysis of microcystin-LR. *Toxicol* 3142: 1-40 .
- Andrinoloa D Sedanb D Teleseb L Aurab C Maserab S Giannuzzia L Marrac CA and Alaniz MJT. 2008. Hepatic recovery after damage produced by sub-chronic intoxication with the cyanotoxin microcystin LR. *Toxicol* 51: 457-67 .
- Billam M Mukhi S Tang L Gao W and Wang J. 2008. Toxic response indicators of microcystin-LR in F344 rats following a single dose treatment. *Toxicol* 3139: 3-4 .
- Chen T Wang Q Cui J Yang W Shi Q Hua Z Ji J and Shen P. 2005. Induction of apoptosis in mouse liver by microcystin-LR. *Mol. & Cell. Proteo.* 4(7): 958-74.
- Dawson RM. 1998. The toxicology of microcystins. *Toxicol* 36 (7): 953-62.
- Denisov ET , Denisova TG and Pokidova TS. 2003. Handbook of free radical initiators, John Wiley & Sons, Inc, Hoboken, New Jersey, USA, 22-57.
- Ding XS Li XY Duan HY Chung IK and Lee JA. 2006. Toxic effects of *Microcystis* cell extracts on the reproductive system of male mice. *Toxicol* 48: 973-9.
- Doyle A and Griffiths JB. 1998. Cell and tissue culture: laboratory procedures in biotechnology. John Wiley & sons, Baffins Lane Chichester, West Sussex P019 1UD, England, 71-4.
- Duan L Carrier DJ. and Clausen EC. 2004. Silymarin extraction from milk thistle using hot water Spring. *App. Biochem. and Biotech.* 59(6): 113-6.
- Dufour JF and Clavien PA. 2005. Signaling Pathways in Liver Diseases, Springer-Verlag, Berlin, Heidelberg, Germany, 47-59.
- Fawell JK Mitchell RE Everett DJ and Hill RE. 1999. The toxicity of cyanobacterial toxins in the mouse: I Microcystin-LR. *Human & Experimental Toxicol.* 18(3): 162-7.
- Gehring MM Downs KS Downing TG Naude RJ and Shephard EG. 2003. An investigation into the effect of selenium supplementation on microcystin hepatotoxicity. *Toxicol* 41: 1-8.
- Gupta US Guha S. 2006. Microcystin toxicity in a freshwater fish, *Heteropneustes fossilis* (Bloch). *Current Science* 91(9): 1261-71.
- Hosseinzadeh H Parvardeh S Asl MN Sadeghnia HR and Ziaee T. 2007. Effect of thymoquinone and nigella sativa seeds oil on lipid peroxidation level during global cerebral ischemia-reperfusion injury in rat hippocampus. *Phytomed.* 14: 621-7.
- Jayaraj R., Anand T. & Lakshmana Rao PV. 2006. Activity and gene expression profile of certain antioxidant enzymes to microcystin-LR induced oxidative stress in mice. *Toxicol.* 220(2,3): 136-46 .
- Joshi G., Sultana R., Tangpong J., Cole MP., Clair D., Vore M., Estus S. and Butterfield DA. 2005. Free radical mediated oxidative stress and toxic side effects in brain induced by the anti cancer drug adriamycin: Insight into chemobrain. *Free Rad. Res.* 39(11): 1147-54.
- Kalapos MP. 2008. The tandem of free radicals and methylglyoxal. *Chemico Biol. Inter.* 171(3): 251-271.
- Kanaze FI., Gabrieli C., Kokkalou E., Georarakis M., Niopas I. 2005. Simultaneous reversed-phase high-performance liquid chromatographic method for the determination of diosmin hesperidin and naringin in different citrus fruit juices and pharmaceutical formulations. *J. Pharm. Biomed Analysis* 33: 243-9.
- Lakshmana Rao PV, Gupta N. and Jayaraj R. 2004. Screening of certain chemoprotectants against cyclic peptide toxin microcystin-LR. *Indian J. Pharmacol.* 36(2): 87-92.
- Lawton, LA Edwards C and Codd GA. 1994. Extraction and high performance liquid chromatographic method for the determination of microcystins in raw and treated waters. *Analyst* 119: 1525-1530.
- Leão JC Geracitano LA Monserrat JM Amado LL and Yunes JS. 2008. Microcystin-induced oxidative stress in *Laeonereis acuta* (Polychaeta, Nereididae). *Marine Environ. Res.* 3192: 1-10 .
- Lohse DL Denu JM and Dixon JE. 1995. Insights derived from the structures of the Ser/Thr phosphatases calcineurin & protein phosphatase. *Structure* 3(10): 987-90.

- Lombardo M Pinto FC Vieira JM Honda RY Carvalho LR & Kiyota S. (2006) Isolation and structural characterization of microcystin - LR and three minor oligopeptides simultaneously produced by *Radiocystis feernandoi* (Chroococcales, cyanobacteria): A Brazilian toxic cyanobacterium. *Toxicon* 47: 560–66.
- Looa AY Jaina K and Darahb I. 2007. Antioxidant and radical scavenging activities of the pyroligneous acid from a mangrove plant, *Rhizophora apiculata*. *Food Chem.* 104(1): 300-7.
- Mackintosh C Beattie KA Klumpp S Cohen P and Codd GA. 1990. Cyanobacterial microcystin-LR is a potent and specific inhibitor of protein phosphatases 1 and 2a from both mammals and higher plants. *Federation of the European Biochemical Society Lett.* 264: 187–92.
- Mereish KA Bunner DL Ragland DR and Creasia DA. 1991. Protection against Microcystin-LR-induced hepatotoxicity by silymarin: Biochemistry, histopathology, and lethality. *Pharma. Res.* 8(2): 273-277.
- Oriola GA and Lawton LA. 2005. Detection & quantification of toxins of *Microcystis aeruginosa* (PCC 7820) by HPLC & protein phosphatase inhibition assay effect of blending various collectors at bulk. *Afri. J. Sci. Technol.* 6(1): 1-10.
- Pinho GL Rosa CM Maciel FE Binachini A Yunes JS Proenca LA and Monserrat JM. 2005. Antioxidant responses and oxidative stress after microcystin exposure in the hepatopancreas of an estuarine crab species. *Ecotoxicol. Environ. Safety* 61: 353-60.
- Rao PVL Jayaraj R and Bhaskar ASB. 2004. Protective efficacy and the recovery profile of certain chemoprotectants against lethal poisoning by microcystin-LR in mice. *Toxicon* 44: 723-730.
- Ratliff DM Vander Jagt DJ Eaton RP and Vander Jagt DL. 1996. Increased levels of methylglyoxal-metabolizing enzymes in mononuclear and polymorphonuclear cells from insulin-dependent diabetic patients with diabetic complications: aldose reductase, glyoxalase I and Glyoxalase II. *J Clin. Endocrinol. Metabol.* 81(2): 488-92.
- Spoof L and Meriluoto J. 2002. Rapid separation of microcystins and nodularin using a monolithic silica C18 column. *J Chromat. A* 947: 237–45.
- Szasz G. (1974) *Methods of Enzymatic Analysis*. Natural Toxins 2: 714-715.
- Weng D Lu Y Wei Y Liu Y and Shen P. 2007. The role of ROS in microcystin-LR- induced hepatocyte apoptosis & liver injury in mice. *Toxicol.* 232: 15-23
- Xu C. Shu W Qiu Z Chen J Zhao Q. and Cao J. 2007. Protective effects of green tea polyphenols against subacute hepatotoxicity induced by microcystin-LR in mice. *Environ. Toxicol. Pharma.* 24: 140-8.
- Yang J Yoneda K Morita E Imamura S Nam K Lee E and Steinert P. 1997. An alanine to proline mutation in the 1a rod domain of the keratin 10 chain in epidermolytic hyperkeratosis. *J. Inves. Dermatol.* 109:692–4.
- Yuan M Carmichael WW and Hilborn ED. 2006. Microcystin analysis in human sera and liver from human fatalities in Caruaru, Brazil. *Toxicon* 48: 627–40.

