

# Estimation of Grape Seed Oil Alleviative Role on Cadmium Toxicity in Male Mice

Osama H. Elhamalawy<sup>1</sup> and Aida I. El makawy<sup>2\*</sup>

<sup>1</sup>Department of Environment and Bio-Agriculture, Faculty of Agriculture, Al-Azhar University, Cairo; <sup>2</sup>Department of Cell Biology, National Research Center, 33 El Bohouth Street, Dokki- Giza-Egypt-P.O.12622.

Received April 14, 2019; Revised May 18, 2019; Accepted May 28, 2019

## Abstract

Cadmium (Cd) is a non-degradable environmental pollutant, and has the ability to collect in the body organs causing humans and animals health risks. The present study was conducted to estimate the protective role of grape seed oil (GSO) against cadmium chloride (CdCl<sub>2</sub>) induced toxicity in male mice. Phenolics content and antioxidant capacity of GSO were measured. Male mice had gavages of CdCl<sub>2</sub> at a dose of (7.48 mg/kg body weight (b. w.) once a day for 30 successive days, whereas they were orally administered GSO at 1 mL/kg b. w. once a day for 30 successive days either pre or plus CdCl<sub>2</sub>. After the treatment, liver enzymes, sperm characteristics and organs weights were investigated. Genotoxicity was evaluated using comet assay and micronucleus test. Results revealed that CdCl<sub>2</sub> induced elevation in ALT and AST, significantly increased the relative weights of liver, decreased the testes weight, sperm count and declined motility and elevated DNA damage in tested cells compared to control group. In contrast, administration of GSO at two regimens restored the organs' weights to the normal range, ameliorated liver enzymes, improved the sperm physical characteristics and alleviated DNA damage compared to CdCl<sub>2</sub> treated mice. Furthermore, GSO pre-administration was the best regimen in the attenuation of the toxic effects of CdCl<sub>2</sub>. It can be concluded that GSO may possess a protective role against the toxicity of CdCl<sub>2</sub> male mice.

**Keywords:** Grape seed oil, Cadmium chloride, Organs weights, Liver enzymes, Sperm physical characters, Genotoxicity.

## 1. Introduction

Heavy metals are a paradigm of pollutants. They are stable high-density metals present in the environment because they are one of the components of earth's crust (Tchounwou *et al.*, 2012). The heavy metals quantity in the air, water, soil and tissues of living organisms has increased because of the human activities that caused their increase in the environment (Gana and Toba, 2015). Heavy metals are toxic, but the dose, degree of exposure and the chemical form of these metals lead to differences in the extent of their toxicity (Tchounwou *et al.*, 2012).

Cadmium (Cd) is a major industrial and environmental pollutant that is principally produced from smelting, mining, electroplating, battery industrialized, paints and plastics. The wide route of exposure to Cd mostly results from smoking, air pollution and Cd contaminated foods and water utilization (Honda *et al.*, 2010). It is accumulated in the liver, kidney and testis, so it is considered as one of the common poisonous heavy metals. Cd causes a broad range of negative health hazards including hepatotoxicity (Kisok, 2012). Wang *et al.* (2017) reported that cadmium is an endocrine disruptor known to apply toxic effects on the testes, with indicated sperm dysfunction during both chronic and acute cadmium

exposure. In addition, Cd is known as human carcinogen that attacks DNA directly and disturbs DNA repair mechanism (Murugavel and Pari, 2010). It is confirmed that Cd induces oxidative stress through the creation of free radicals that may damage protein, lipid, enzymes and DNA (Naskar *et al.*, 2010).

Antioxidants provide potent free radicals scavengers and thwart the incidence of disease (Islam *et al.*, 2011). In recent times, using natural foodstuffs for diseases handling has been broadly suggested due to their protective effects (Soliman *et al.*, 2013; 2015). Such findings suggest that it is necessary to identify alternative drugs for protection against toxicity and the treatment of human diseases. Grape (*Vitis vinifera*) is one of the world's major fruit crops, and grape seed extract is a multifarious matrix with approximately 40% fiber, 16% oil, 11% proteins, and 7% phenols such as tannins and mineral (Shi *et al.*, 2003). The grape seed oil (GSO) has a lot of uses for food additive, cosmetics, controlling diseases and wound healing (Shivananda *et al.*, 2011). Many researchers have mentioned plentiful health benefits of GSO, where it is considered as a potent antioxidant for its polyphenols, flavonoids, unsaturated fatty acids and vitamin E contents (Hassanein and Abedel-Razek, 2009; Kikalishvili *et al.*, 2011; Hasseeb *et al.*, 2013). Grape seed extracts (GSE)

\* Corresponding author e-mail: aelmakawy@yahoo.com.

possess antimutagenic and anticarcinogenic properties by inhibiting enzymes of free radicals productions (Maier *et al.*, 2009). Moreover, the exploitation of GSE may be helpful in withdrawing the side effects of chemotherapeutic agents (Aysun *et al.*, 2008). In addition, it is recognized that the GSO could supply protection against free radicals induced oxidative stress and cellular damage caused after exposure to heavy metals (Shinagawa *et al.*, 2015).

Consequently, the aim of this study was to estimate the potential protective role of grape seed oil against the toxicity induced by cadmium in liver and testis of mice. The study was carried out by monitoring DNA damage by micronucleus test and comet assay in male mice, by examining liver function (enzymes), sperm analysis, and by determining the body weight of animals.

## 2. Materials and Methods

### 2.1. Chemicals

Cadmium chloride (CdCl<sub>2</sub>) was purchased from Sigma-Aldrich Company (St. Louis, MO, USA). Grape seed commercial oil (GSO) was purchased from EL Captin Company (Al Obour City, Cairo, Egypt). The aspartate aminotransferase (AST) and alanine aminotransferase (ALT) kits were purchased from Biodiagnostic Co., Egypt. All other chemicals were of analytical grade and purchased from standard commercial suppliers.

### 2.2. Animals and treatments

Male Swiss albino mice weighing 24±5 g (10-12 week old) were purchased from the Theodor Bilharz Research Institute, Giza, Egypt. Animals were housed in polypropylene cages (43cm× 30cm × 15cm, five mice per cage) with stainless steel covers in the Animal House of Environment and Bio-agriculture Department, Faculty of Agriculture, Al-Azhar University. Animals were kept under controlled temperature (23±4 °C), 50–55% relative humidity and photoperiod of 12 h light:12 h dark cycle. Throughout the experimental period, animals were maintained on standard mice pellet food (Salaam Feed Factory, El marg, Cairo) and tap water *ad libitum*. The mice were allowed to adapt to their surrounding environment for 2 weeks prior to start the experiments. All animals received human care in compliance with the guidelines of the Animal Care and Use Committee of the National Institutes of Health (NIH publication 86-23 revised 1985).

After acclimation, the animals were randomly divided into five groups (n = 10) according to approximately equal mean body weight (b.w.) and administered orally for 30 successive days with CdCl<sub>2</sub> (1/25 LD<sub>50</sub>, 7.48 mg/kg b.w.) according to Yang *et al.* (2012a) and/or GSO (1mL/kg b.w.) according to Mokhtari *et al.* (2011). Experimental groups were as follows: Control group: animals were orally administered with saline. GSO group: animals were orally administered with GSO at dose of (1mL/kg b.w.). CdCl<sub>2</sub> group: animals were orally administered with CdCl<sub>2</sub> at dose of (1/25 LD<sub>50</sub>). GSO pre-CdCl<sub>2</sub> group: animals were orally administered with 1 mL/kg b.w. GSO then with CdCl<sub>2</sub> (1/25 LD<sub>50</sub>). GSO plus CdCl<sub>2</sub> group: animals were orally administered with 1 mL/kg b.w. GSO plus CdCl<sub>2</sub> (1/25 LD<sub>50</sub>).

### 2.3. Grape seed oil total phenolic content

Total phenolic content (TPC) was determined using the Folin–Ciocalteu's reagent according to the method reported by Lin and Tang (2007) at 760 nm with a spectrophotometer (UV-1601; Shimadzu, Tokyo, Japan), and the quantification was done on the basis of the standard curve of gallic acid concentration ranging between 10 to 80 mg/mL (r<sup>2</sup>= 0.99).

### 2.4. Measurement of antioxidant activity of grape seed oil

The ability of GSO at 50, 100 and 200µL to scavenge 2.9 mL of 1, 1'-diphenyl 1-2-picrylhydrazyl (DPPH) free radical was estimated by the method of Singh *et al.* (2002).

### 2.5. Relative organs weights

At the termination of the experiments, internal organs such as liver and testes were dissected out, trimmed of excess fat and weighted. The organs weight was presented as relative organ weight as follows:

$$\text{Relative organ weight} = \frac{\text{Organ weight (g)}}{\text{Final body weight (g)}} \times 100$$

### 2.6. Liver function investigation

Aspartate aminotransferase (AST) and alanine aminotransferase (ALT) liver enzymes activities were measured in serum by the method of Reitman and Frankel (1957).

### 2.7. Sperm collection and analysis

The epididymides from each mouse were removed and sperm was collected as quickly as possible after dissection. Epididymides was excised and minced in 1 mL of phosphate buffered saline (pH 7.2) to obtain sperm suspension; the sperms were filtered afterwards through a nylon mesh. The sperm count was assessed from right cauda epididymides while sperm motility was analyzed from the left one. Sperm count was performed according to Narayana *et al.* (2002) using a Neubauer hemocytometric chamber. Sperm motility was assessed by counting the motile sperms. Approximately 10 µL of sperm suspension was layered onto a warmed microscope slide. In each semen sample, at least 10 microscopic fields were examined with at least 100 sperm/field counted. The percentage of motile sperm was determined as described by Kvist and Bjorndahl (2002).

### 2.8. Bone marrow micronucleus assay

Bone marrow sampling, preparations of slide and scoring of micronucleated polychromatic erythrocytes (MNPCE) were done as described by Hayes *et al.* (2009). In brief, bone marrow was gently flushed out in fetal calf serum, centrifuged at 1000 rpm for 5 min and the medium was decanted. Pellet obtained was dispersed in 0.25 mL fetal calf serum, smeared on clean slides, fixed in 70% methanol, air dried and stained with May-Grunwald/Giemsa protocol. Minimum of 2000 polychromatic erythrocytes/mouse were scored from treated or control group.

### 2.9. Single cell gel Electrophoresis in liver and testes

Single cell gel Electrophoresis (comet assay) was performed in liver and testes cells according to Bandyopadhyay *et al.* (2008). Briefly, 50 µL of cell suspension was mixed with 100 µL of 1 % low melting point (LMP) agarose and added to fully frosted slides

coated with 80  $\mu\text{L}$  of 1 % normal melting point (NMP) agarose. The cells were then incubated in a lysis solution (2.5 mol  $\text{L}^{-1}$  NaCl, 100 mmol  $\text{L}^{-1}$  EDTA, 10 mmol  $\text{L}^{-1}$  Tris-HCL, 1 % Triton X-100, pH 10) at 4 °C for at least 2 h, at which the slides were placed into an alkaline solution (300 mmol  $\text{L}^{-1}$  NaOH, 1 mmol  $\text{L}^{-1}$  EDTA, pH 13) at 4 °C for 20 min so as to allow DNA unwinding, and electrophoresed at 25 V (300 mA) for 20 min. Finally, the slides were neutralized in 400 mmol/L Tris buffer (pH 7.5) for 15 min and stained with ethidium bromide (5  $\mu\text{g mL}^{-1}$ ). Images of 50 randomly selected nuclei per experimental group were captured using a fluorescence microscope (Eclipse 800, Nikon, Tokyo, Japan) and analyzed with image analysis software (Comet Assay IV, Perceptive Instruments, Suffolk, UK). Scored parameters included tail length, DNA percentage in tail and olive tail moment (OTM). Tail length is the maximum distance that the damaged DNA migrates from the center of the cell nucleus. DNA percentage in tail is the DNA content that migrates from the nucleus into the comet tail. OTM is the product of the tail length and percentage DNA, which gives a more integrated measurement of overall DNA damage in the cell.

### 2.10. Statistical analyses

Statistical analyses were performed with SPSS 16 software. Experimental data was analyzed using one-way analysis of variance (ANOVA) followed by Duncan's post hoc test for multiple comparisons between pairs. All values were expressed as mean  $\pm$  S.D. and differences were considered as significant when ( $P \leq 0.05$ ).


## 3. Results

### 3.1. Oil analysis

GSO was found to have high level of phenolic content (42.17 $\pm$ 0.55 mg gallic acid /100g oil) and antioxidant activity (84.35 $\pm$ 0.35) as show in Table 1. In addition, antioxidant activity analysis of GSO revealed that its scavenging activity was raised gradually with increasing the concentration as shown in Figure 1.

**Table 1.** Phenolic content and antioxidant activity in grape seed oil.

Character	M $\pm$ SD
Phenolic content (mg gallic acid /100g oil)	42.17 $\pm$ 0.55
Antioxidant activity %	84.35 $\pm$ 0.35



**Figure 1.** Grape seed oil antioxidant activity.

### 3.2. Relative organs weights

According to data presented in Table 2, there was no significant difference between both liver and testes relative weights of control and GSO treated groups. Oral administration of  $\text{CdCl}_2$  to male mice for 30 successive days significantly increased ( $P \leq 0.05$ ) the relative weights of liver (4.18 $\pm$ 0.05) and decreased the testes weight (0.65 $\pm$ 0.03) as compared to control (3.39 $\pm$ 0.06 and 0.83 $\pm$ 0.03), respectively. On the other hand, GSO administration either pre or plus  $\text{CdCl}_2$  induced a significant decrease ( $P \leq 0.05$ ) in the relative weights of liver (3.39 $\pm$ 0.04 and 3.33 $\pm$ 0.07, respectively) and increase in testes (0.77 $\pm$ 0.02 and 0.74 $\pm$ 0.02, respectively) compared with  $\text{CdCl}_2$  treated mice.

**Table 2.** Relative organs weights of treated male mice with cadmium chloride ( $\text{CdCl}_2$ ) and / or grape seed oil (GSO) for 30 consecutive days.

Treatments	Relative organs weights (g)	
	Liver	Testes
Control	3.39 $\pm$ 0.06 <sup>bc</sup>	0.83 $\pm$ 0.03 <sup>a</sup>
GSO (1 mL/kg b.wt)	3.46 $\pm$ 0.03 <sup>b</sup>	0.86 $\pm$ 0.02 <sup>a</sup>
$\text{CdCl}_2$ (1/25 LD <sub>50</sub> )	4.18 $\pm$ 0.05 <sup>a</sup>	0.65 $\pm$ 0.03 <sup>c</sup>
GSO pre- $\text{CdCl}_2$	3.39 $\pm$ 0.04 <sup>bc</sup>	0.77 $\pm$ 0.02 <sup>b</sup>
GSO plus $\text{CdCl}_2$	3.33 $\pm$ 0.07 <sup>b</sup>	0.74 $\pm$ 0.02 <sup>b</sup>

Data are expressed as means  $\pm$  SD. Mean values in the same column within each parameter bearing the same superscript do not differ significantly ( $P \leq 0.05$ ).

### 3.3. Liver function

The mean values of serum liver enzymes in  $\text{CdCl}_2$  and/or GSO treated male mice are illustrated in Table 3. A significant increase ( $P \leq 0.05$ ) in the mean values of ALT (43.50 $\pm$ 3.31) and AST (46.25 $\pm$ 4.11) was observed following the treatment with  $\text{CdCl}_2$  for 30 successive days compared with control (24.50 $\pm$ 1.29 and 31.00 $\pm$ 1.83), respectively. In contrast, oral administration of GSO either pre or plus  $\text{CdCl}_2$  produced a significant decrease ( $P \leq 0.05$ ) in serum content of ALT (30.75 $\pm$ 2.06 and 34.25 $\pm$ 3.30, respectively) and AST (34.25 $\pm$ 2.50 and 38.25 $\pm$ 3.59, respectively) than that of  $\text{CdCl}_2$  treated male mice.

**Table 3.** Serum liver enzymes of cadmium chloride ( $\text{CdCl}_2$ ) and / or grape seed oil (GSO) treated male mice.


Treatments	ALT (U/L)	AST (U/L)
Control	24.50 $\pm$ 1.29 <sup>c</sup>	31.00 $\pm$ 1.83 <sup>c</sup>
GSO (1 mL/kg bw)	23.50 $\pm$ 1.73 <sup>c</sup>	29.75 $\pm$ 2.22 <sup>c</sup>
$\text{CdCl}_2$ (1/25 LD <sub>50</sub> )	43.50 $\pm$ 3.31 <sup>a</sup>	46.25 $\pm$ 4.11 <sup>a</sup>
GSO pre- $\text{CdCl}_2$	30.75 $\pm$ 2.06 <sup>b</sup>	34.25 $\pm$ 2.50 <sup>bc</sup>
GSO plus $\text{CdCl}_2$	34.25 $\pm$ 3.30 <sup>b</sup>	38.25 $\pm$ 3.59 <sup>b</sup>

Data is expressed as means  $\pm$  SD. Mean values in the same column within each parameter bearing the same superscript do not differ significantly ( $P \leq 0.05$ ).

### 3.4. Sperm analysis

As shown in Figure 2, treatment with  $\text{CdCl}_2$  for 30 successive days caused a significant decline ( $P \leq 0.05$ ) in the mean values of sperm count (284.20 $\pm$ 29.51) and motility (54.00 $\pm$ 5.09) compared with control

(465.40±22.41 and 85.00±3.53), respectively. By contrast, supplementation of GSO to male mice either before or with CdCl<sub>2</sub> significantly improved ( $P \leq 0.05$ ) the decrease of sperm count (447.20±22.70 and 400.60±28.02, respectively) and motility (82.00±2.55 and 76.25±2.44, respectively) compared with CdCl<sub>2</sub> treated mice.



**Figure 2.** Sperm count and motility in cadmium chloride (CdCl<sub>2</sub>) and/or grape seed oil (GSO) treated mice for 30 consecutive days. Data is expressed as means ± SD. Mean values in the same column within each parameter bearing the same superscript do not differ significantly ( $P \leq 0.05$ ).

**Table 4.** Frequencies of PCEs, NCEs, MN and PCEs/NCEs percentage in bone marrow cells of cadmium chloride (CdCl<sub>2</sub>) and/or grape seed oil (GSO) treated mice.

Treatments	PCEs /2000Cell	NCEs /2000Cell	MNPCEs /2000Cell	PCEs/NCEs %
Control	1440.67±32.44 <sup>b</sup>	559.33±32.44 <sup>c</sup>	5.33±1.33 <sup>c</sup>	0.025±0.002 <sup>b</sup>
GSO (1 mL/kg b.w.)	1544.00±16.77 <sup>a</sup>	456.00±16.77 <sup>d</sup>	4.00±1.15 <sup>c</sup>	0.033±0.001 <sup>a</sup>
CdCl <sub>2</sub> (1/25 LD <sub>50</sub> )	1181.33±32.91 <sup>d</sup>	818.67±32.91 <sup>a</sup>	40.67±6.76 <sup>a</sup>	0.014±0.001 <sup>d</sup>
GSO pre-CdCl <sub>2</sub>	1384.67±27.55 <sup>bc</sup>	615.33±27.55 <sup>bc</sup>	16.67±2.90 <sup>b</sup>	0.022±0.002 <sup>bc</sup>
GSO plus CdCl <sub>2</sub>	1311.33±28.48 <sup>c</sup>	688.67±28.48 <sup>b</sup>	26.00±2.00 <sup>b</sup>	0.018±0.001 <sup>cd</sup>

Data is expressed as means ± SD. Mean values in the same column within each parameter bearing the same superscript do not differ significantly ( $P \leq 0.05$ ). \* PCEs: Polychromatic erythrocytes NCEs: Normochromatic erythrocytes MN: Micronuclei.

### 3.6. Effect of CdCl<sub>2</sub> and/or GSO on tailed cells, tail length, tail DNA % and olive tail moment in liver and testes

The mean values of tailed, intact cells, tail length, tail DNA % and olive tail moment in CdCl<sub>2</sub> and/or GSO treated liver and testes are presented in Tables 5 and 6. There were no significant differences in all tested parameters between control and GSO treated animals in both organs cells. The values of tailed cell percentage, tail length, DNA percentage in tail and olive tail moment in liver cells of male mice treated with CdCl<sub>2</sub> were

**Table 5.** The mean values of intact, tailed cells, tail length, tail DNA % and olive tail moment in cadmium chloride (CdCl<sub>2</sub>) and/or grape seed oil (GSO) treated liver.

Treatments	Intact cells (%)	Tailed cells (%)	Tail length (µm)	Tail DNA (%)	Olive tail moment (µm)
Control	86.17±0.38 <sup>ab</sup>	13.83±0.38 <sup>cd</sup>	9.04±0.57 <sup>d</sup>	13.04±0.37 <sup>d</sup>	1.18±0.07 <sup>d</sup>
GSO (1 mL/kg b.w.)	87.30±0.79 <sup>a</sup>	12.70±0.79 <sup>d</sup>	8.44±0.46 <sup>d</sup>	12.22±0.73 <sup>d</sup>	1.03±0.04 <sup>d</sup>
CdCl <sub>2</sub> (1/25 LD <sub>50</sub> )	70.33±3.26 <sup>d</sup>	29.67±3.26 <sup>a</sup>	19.56±0.58 <sup>a</sup>	21.78±0.41 <sup>a</sup>	4.26±0.09 <sup>a</sup>
GSO pre-CdCl <sub>2</sub>	83.40±1.44 <sup>bc</sup>	16.60±1.44 <sup>bc</sup>	11.40±0.77 <sup>c</sup>	14.53±0.55 <sup>c</sup>	1.66±0.14 <sup>c</sup>
GSO plus CdCl <sub>2</sub>	80.73±1.86 <sup>c</sup>	19.27±1.86 <sup>b</sup>	13.40±0.61 <sup>b</sup>	16.16±0.31 <sup>b</sup>	2.16±0.10 <sup>b</sup>

Data is expressed as means ± SD. Mean values in the same column within each parameter bearing the same superscript do not differ significantly ( $P \leq 0.05$ ).

### 3.5. Effect of CdCl<sub>2</sub> and/or GSO on micronuclei number and PCEs/NCEs percentage in bone marrow

Results in Table 4 showed that there were no statistically significant differences in the numbers of MN in the control and GSO-treated groups. While, CdCl<sub>2</sub> significantly increased ( $P \leq 0.05$ ) the frequency of MNPCEs (40.67±6.76) as compared to control (5.33±1.33). On the other side, administration of GSO either pre or plus CdCl<sub>2</sub> decreased significantly ( $P \leq 0.05$ ) the frequencies of MNPCEs (16.67±2.90 and 26.00±2.00) than in CdCl<sub>2</sub> group (40.67±6.76). Concerning to the PCEs/NCEs percentage, data analysis revealed that CdCl<sub>2</sub> caused marked toxicity in bone marrow cells, where it is significantly decreased ( $P \leq 0.05$ ) the percentage of PCEs/NCEs. Along with that, GSO ameliorated the toxicity of CdCl<sub>2</sub> by increasing the PCEs/NCEs percentage.

significantly increased ( $P \leq 0.05$ ) compared to those of control. However, supplementation of GSO to male mice either pre or plus CdCl<sub>2</sub> caused a significant alleviation ( $P \leq 0.05$ ) in the values of above-mentioned parameters than in CdCl<sub>2</sub> treated animals values. In addition, CdCl<sub>2</sub> caused a significant rise ( $P \leq 0.05$ ) in all values of testes cells compared to control. By contrast, administration of GSO plus CdCl<sub>2</sub> with both patterns repressed significantly ( $P \leq 0.05$ ) the increase in the all previously mentioned parameters rate as compared to those caused by CdCl<sub>2</sub>.

**Table 6.** Effect of cadmium chloride (CdCl<sub>2</sub>) and/or grape seed oil (GSO) on Tailed cells, Tail length, Tail DNA % and Olive tail moment in testes.

Treatments	Intact cells(%)	Tailed cells (%)	Tail length (µm)	Tail DNA (%)	Olive tail moment (µm)
Control	87.23±0.65 <sup>a</sup>	12.77±0.65 <sup>d</sup>	7.54±0.58 <sup>cd</sup>	10.61±0.62 <sup>d</sup>	0.79±0.02 <sup>d</sup>
GSO	87.70±0.36 <sup>a</sup>	12.30±0.36 <sup>d</sup>	6.91±0.59 <sup>d</sup>	10.49±0.48 <sup>d</sup>	0.72±0.05 <sup>d</sup>
CdCl <sub>2</sub>	67.76±1.35 <sup>d</sup>	32.24±1.35 <sup>a</sup>	14.90±0.89 <sup>a</sup>	18.40±0.65 <sup>a</sup>	2.74±0.19 <sup>a</sup>
GSO pre-CdCl <sub>2</sub>	82.43±1.02 <sup>b</sup>	17.57±1.02 <sup>c</sup>	8.61±0.54 <sup>c</sup>	12.15±0.51 <sup>c</sup>	1.04±0.03 <sup>c</sup>
GSO plus CdCl <sub>2</sub>	77.86±2.04 <sup>c</sup>	22.14±2.04 <sup>b</sup>	9.87±0.48 <sup>b</sup>	14.45±0.53 <sup>b</sup>	1.43±0.11 <sup>b</sup>

Data is expressed as means ± SD. Mean values in the same column within each parameter bearing the same superscript do not differ significantly ( $P \leq 0.05$ ).

#### 4. Discussion

Grape seed products are nutraceutical agents generally utilized as health dietary supplements. The concern about GSO as a functional food has increased because of its high content of phenolic compounds, vitamin E, unsaturated fatty acids (UFAs) and phytosterols (Karaman *et al.*, 2015). This was in line with the result of oil analysis, which confirmed that GSO possess a high antioxidant capacity due to high content of gallic acid, catechin, proanthocyanidins, procyanidins and epicatechin.

Cadmium intoxicated mice showed clear increase in liver weight, may be because the hypertrophy was induced in the liver because of cadmium toxicity. This result is consistent with a previous report indicating that CdCl<sub>2</sub> toxicity leads to increasing the weight of liver in rats (Bashir *et al.*, 2014). Otherwise, cadmium produced loss in testis weight. This was consistent with the Elgawish and Ghanem (2014) study; they indicated that the reproductive organs weight was decreased in rats exposed to CdCl<sub>2</sub> as compared to the control. The decline in testes and epididymis weight may be ascribed to the inhibition of spermatogenesis, decreased elongated spermatids and steroidogenic enzyme activity (Takahashi and Oishi, 2003). Salem and Salem, (2016) observed that cadmium induced liver weight elevations and testes weights reduction. Conversely, GSO restored the organs weight to the normal range. Our results match with Bashir *et al.* (2014); they found that cadmium decreased water and food intake followed with retardation in growth rate and increase in liver weight. Authors interpreted that Cd accumulation causes disturbances in the body and liver weight of rats due to tissue damage and reduction in their functions. All these alterations induced by Cd intoxication were significantly restored to near normal levels upon pre-administration of GSE. In our study, marked hepatic damage was observed because of the significant elevation of the serum hepatic enzymes (ALT, AST) in the CdCl<sub>2</sub> group. These characteristic features of CdCl<sub>2</sub>-induced liver toxicity are comparable to those formerly reported by other investigators (Obioha *et al.*, 2009; Renugadevi and Prabu, 2010; Shati, 2011; Salem and Salem, 2016). The elevation in liver enzymes activities in CdCl<sub>2</sub> intoxicated animals had been attributed to cellular leakage in the organ and loss of functional integrity of membrane architecture of hepatocytes (Salem and Salem, 2016). On the other hand, the disturbance in the levels of the hepatic enzymes in CdCl<sub>2</sub> intoxicated mice was ameliorated by both pre- and co-administration with the GSO, as there was a significant

decrease in the levels of these enzymes in mice treated with GSO pre/plus CdCl<sub>2</sub>. This was in agreement with Dogan and Celik (2012) as they indicated that the grape seed extract (GSE) efficiently protected the rats against alcohol-induced hepatotoxicity, as evidenced by AST, ALT enzyme levels reduction. In addition, Abdul-Hamid *et al.* (2018) found that GSE administration with amiodarone revealed obvious improvement for liver enzyme activity compared to amiodarone-treated animals. Bishayee *et al.* (2010) suggested that polyphenol existing in grape seed could reduce hepatic enzymes through the antioxidant actions.

Our findings clarified that CdCl<sub>2</sub> administration caused significant decrease in sperm count and motility. These findings were in consent with those of earlier studies by Oliveira *et al.* (2009) and Kaur and Sharma (2015) who observed significant reduction in sperm physical characters in CdCl<sub>2</sub> exposed mice. Acharya *et al.* (2008) clarified that reactive oxygen species (ROS) generated by cadmium can cause spermatozoa lipid peroxidation and deleterious effects on cell membrane phosphatides, inducing sperm DNA oxidation and consequently sperm abnormalities and decline in alive sperm count. The GSO caused amelioration of physical properties of semen in mice. This was in agreement with Al-Shahari and El-kott (2019); they demonstrated a defensive impact of grape seeds extract against the poisoning of monosodium glutamate, whereas Yildirim *et al.* (2011) indicated that the GSE affects sperm movement sperms number, and motility (%), with less immotile and abnormal sperm morphology when contrasted with control. Also, Al-Saeed (2016) reported that GSO caused sperm count elevation in the diazinon treated rats. This could be due to the capability of GSO to either interfere with the spermatogenic processes in the seminiferous tubules, epididymal functions of testosterone on hypothalamic release factor and anterior pituitary secretion of gonadotropins that may lead to activation of spermatogenesis.

Our data showed that CdCl<sub>2</sub> induced DNA damage measured by MN and comet assays in liver and testes cells which is considered as indicator of its genotoxicity. Likewise, Abd-El-Moneim *et al.* (2017); El-Habit and Abdel Moneim (2014) reported significant elevations in micronuclei number in murine bone marrow cells exposed to CdCl<sub>2</sub>. Moreover, Mahrous *et al.* (2015) discovered highly significant frequencies of micronuclei in Nile tilapia fish treated with CdCl<sub>2</sub>. Also, the results of DNA damage measured by comet assay were in consent with several studies, like Karimi *et al.* (2012) who observed significant alterations in DNA in mice kidney cells

exposed to CdCl<sub>2</sub> as compared with control mice. Abd-El-Moneim *et al.* (2017) indicated that the rates of DNA damage in liver cells of CdCl<sub>2</sub> contaminated diet fed mice were significantly increased, with significant elevation in comet parameters including (tailed cells %, tail length, tail DNA % and tail moment). In addition, Skipper *et al.* (2016) found significant elevation of DNA damage measured by comet assay in liver carcinoma cells of humans treated with CdCl<sub>2</sub>. Many studies revealed that CdCl<sub>2</sub> oral administration in drinking water of rats enhanced oxidative stress and subsequently the excess of ROS, which may induce DNA damage. Literature data pointed that cadmium induced DNA damage, thus leading to the creation of single and double strand breaks (Yang *et al.*, 2012b; Virk *et al.*, 2013; De Souza Predes *et al.*, 2014). Administration of GSO before or during the treatment with CdCl<sub>2</sub> ameliorates the genotoxicity induced by CdCl<sub>2</sub> in both tested organs. These results were in agreement with study of Cavusoglu *et al.* (2014), which confirmed that oral gavages with grape seed extract significantly ameliorated the indices of hepatotoxicity, nephrotoxicity, lipid peroxidation, and genotoxicity induced by benzene. They showed a significant reduction in the frequency of MNs when compared with the group treated with benzene alone.

## 5. Conclusion

Grape seed oil is wealthy in phenolics, fatty acids and vitamins, with economic value to pharmaceutical and food industry. It has useful properties for health, such as anti-inflammatory, cardioprotective, antimicrobial, and anti-cancer activities, and may interact with cellular and molecular pathways. GSO showed marked alleviative role against CdCl<sub>2</sub> hepatotoxicity and genotoxicity in male mice. These activities could be attributed to high content of phenolics and antioxidant activity which is reflected in renewed liver enzymes, semen character and reduction of DNA damage almost to normal range. Therefore, the GSO may be efficient to diminish the heavy metals health risk.

## References

- Abd-El-Moneim MO, Abd El-Kader AH, Abd El-Rahim HA, Radwan AH, Fadel M and Farag MI. 2017. Modulatory role of *Saccharomyces cerevisiae* against cadmium-induced genotoxicity in mice. *J Arab Soc Med Res.*, **12**: 27-38.
- Abdou MH and Wahb MM. 2016. Neuroprotection of grape seed extract and pyridoxine against triton-induced neurotoxicity. *Oxid Med Cell Longev.*, **5**: 1-8.
- Abdul-Hamid M, Galaly RS, Mahmoud H and Mostafa F. 2018. The protective effect of grape seed and *Ginkgo biloba* against hepatotoxicity induced by the antidysrhythmic drug "amiodarone" in male albino rats. *J Basic Appl Sci.*, **7**: 223-230.
- Acharya UR, Mishra M, Patro J and Panda MK. 2008. Effect of vitamins C and E on spermatogenesis in mice exposed to cadmium. *Reprod Toxicol.*, **25**: 84-88.
- Al- Attar SM. 2017. Antimutagenic effect of grape seed extracted oil on diazinon induced genotoxicity in albino mice. *Iraqi J Cancer Med Genet.*, **10**(1): 56-62.
- AL-Saeed HM. 2016. Effect of grape seed oil on alterations of hematological parameters and semen fluid quality induced by diazinon in male rats. *Indo - Asian J Multidisciplinary Res (IAJMR)*, **2**(6): 843-855.
- Al-Shahari AE and El-kott FA. 2019. Potential effect of grape seeds extract against monosodium glutamate induced infertility in rats. *Int J Pharmacol.*, **15**: 287-294.
- Aysun C, Leylagul K, Ismail K, Sibel H, Recep S, Ahmet O, Okan O and Osman S. 2008. The effect of grape seed extract on radiation- induced oxidative stress in the rat liver. *Turk J Gastroenterol.*, **19**(2): 92-98.
- Bandyopadhyay A, Ghoshal S and Mukherjee A. 2008. Genotoxicity testing of low-calorie sweeteners: aspartame, acesulfame-K, and saccharin. *Drug Chem Toxicol.*, **31**(4): 447-57.
- Bashir N, Manoharan V and Prabu MS. 2014. Protective role of grape seed proanthocyanidins against cadmium induced hepatic dysfunction in rats. *Toxicol Res.*, **3**: 131-141.
- Bishayee A, Darvesh SA, Politis T, and McGory R. 2010. Resveratrol and liver disease: from bench to bedside and community. *Liver Int.*, **30**(8): 1103-1114.
- Cavusoglu K, Yalcin E, Yapar K, Gur B and Cicek F. 2014. The protective role of grape seed extract against chronic toxicity of benzene in Swiss albino mice. *Cumhuriyet Univ Facul Sci J.*, **35**(1): 1-11.
- De Souza Predes F, da Silva Diamante AM, Foglio AM, Camargo AC and Aoyama H. 2014. Hepatoprotective effect of *Arctium lappa* root extract on cadmium toxicity in adult Wistar rats. *Biol Trace Element Res.*, **160**: 250-257.
- Dogan A and Celik I. 2012. Hepatoprotective and antioxidant activities of grape seeds against ethanol-induced oxidative stress in rats. *Br J Nutr.*, **107**(1): 45-51.
- Elgawish RAR and Ghanem ME. 2014. Effect of long term cadmium chloride exposure on testicular function in male albino rats. *Am J Anim Vet Sci.*, **9**: 182-188.
- El-Habit OH and Abdel Moneim AE. 2014. Testing the genotoxicity, cytotoxicity and oxidative stress of cadmium and nickel and their additive effect in male mice. *Biol Trace Elem Res.*, **159**: 364-372.
- Gana AJ and Toba AP. 2015. Environmental pollution and sustainability. *J Res Environ Sci Toxicol.*, **4**(1): 1-9.
- Hassanein M M and Abedel-Razek AG. 2009. Chromatographic quantitation of some bioactive minor components in oils of wheat germ and grape seeds produced as by-products. *J Oleo Sci.*, **58**: 227-233.
- Hasseeb MM, Al-Hizab FA and Hamouda MA. 2013. Impacts of grape seed oil supplementation against the acrylamide induced lesions in male genital organs of rats. *Pak Vet J.*, **33**: 282-286.
- Hayes J, Doherty AT, Adkins DJ, Oldman K and O'Donovan MR. 2009. The rat bone marrow micronucleus test-study design and statistical power. *Mutagenesis* **24** (5): 419-24.
- Honda R, Swaddiwudhipong W, Nishijo M, Mahasakpan P, Teeyakasem W and Ruangyuttikarn W. 2010. Cadmium induced renal dysfunction among residents of rice farming area downstream from a zinc-mineralized belt in Thailand. *Toxicol Lett.*, **198**: 26-32.
- Islam MR, Shahnaj MP, Raihan MO, Hasan SMR and Islam ME. 2011. *In vitro* and *in vivo* antioxidant potential of ethanolic extract of *Syzygium jambos* (L) bark. *Int J Res Ayurveda Pharm.*, **2**: 810-815.
- Karaman S, Karasu S and Tornuk F. 2015. Recovery potential of cold press byproducts obtained from the edible oil industry: physicochemical, bioactive, and antimicrobial properties. *J Agric Food Chem.*, **63**(8): 2305-2313.
- Karimi MM, Sani MJ, Mahmudabadi AZ, Sani AJ and Khatibi SR. 2012. Effect of acute toxicity of cadmium in mice kidney cells. *Iran J Toxicol.*, **6**: 691-698.

- Kaur S and Sharma S. 2015. Evaluation of toxic effect of cadmium on sperm count, sperm motility and sperm abnormality in albino mice. *Int J Adv Res.*, **3**: 335–343.
- Kikalishvili B, Zurabashvili D, Nikolaishvili M, Zurabashvili Z and Giorgobiani I. 2011. The most biological important constances of Rkatsiteli grape oil and its effect as a 5% and 10% food-additive. *Georgian Med News.*, **195**: 85–87
- Kisok K. 2012. Blood cadmium concentration and lipid profile in Korean adults. *Environ Res.*, **112**: 225–229.
- Kvist U and Björndahl L. 2002. **Manual on Basic Semen Analysis**. ESHRE Monographs. Oxford, United Kingdom: Oxford University Press.
- Lin J and Tang C. 2007. Determination of total phenolic and flavonoid contents in selected fruits and vegetables, as well as their stimulatory effects on mouse splenocyte proliferatio. *Food Chem.*, **101**(1): 140-147.
- Mahrous KF, Hassan AM, Radwan HA and Mahmoud MA. 2015. Inhibition of cadmium induced genotoxicity and histopathological changes in Nile tilapia fish by Egyptian and Tunisian montmorillonite clay. *Ecotoxicol Environ Saf.*, **119**: 140–147.
- Maier T, Schieber A, Kammerer D and Carle R. 2009. Residues of grape (*Vitis vinifera* L.) seed oil production as a valuable source of phenolic antioxidants. *Food Chem.*, **112**: 551-559.
- Miltonprabu N, Basher V and Manoharan E. 2016. Hepatoprotective effect of grape seed proanthocyanidins on cadmium-induced hepatic injury in rats: possible involvement of mitochondrial dysfunction, inflammation and apoptosis. *J Toxicol Rep.*, **3**: 63-70.
- Mokhtari M, sarkaki A, Sharifi E and Basiryani E. 2011. Antinociceptive effects of grape seed oil with use of formalin test in male rats. Inter Conference on Food Engineering and Biotechnol (**ICFEB**), 48-53.
- Murugavel P and Pari L. 2010. Effect of diallyl tetrasulfide on cadmium induced oxidative damage in the liver of rats. *Hum Exp Toxicol.*, **26**: 527–534.
- Narayana K, D'Souza UJA and Rao KPS. 2002. Ribavirin induced sperm shape abnormalities in Wistar rats. *Mutat Res.*, **513**: 193–196.
- Naskar S, Islam A, Mazumder UK, Saha P, Haldar PK and Gupta M. 2010. *In vitro* and *in vivo* antioxidant potential of hydro-methanolic extract of *Phoenix dactylifera* fruits. *J Sci Res.*, **2**: 144-157.
- Obioha UE, Suru SM, Ola-Mudathir KF and Faremi TY. 2009. Hepatoprotective potentials of onion and garlic extracts on cadmium-induced oxidative damage in rats. *Biol Trace Elem Res.*, **129**: 143-156.
- Oliveira H, Spanò M, Santos C and Pereira M. 2009. Adverse effects of cadmium exposure on mouse sperm. *Reprod Toxicol.*, **28**: 550–555.
- Reitman A and Frankel S. 1957. A colorimetric method for the determination of serum glutamic oxalacetic and glutamic pyruvic transaminases. *Am J Clin Path.*, **28**: 56.
- Renugadevi J and Prabu SM. 2010. Cadmium-induced hepatotoxicity in rats and the protective effect of naringenin. *Exp Toxicol Pathol.*, **62**: 171-181.
- Salem NA and Salem EA. 2016. Hepatorenal and testicular protective effects of lycopene against cadmium induced toxicity in male rats. *J Nephrol Ther.*, **6**: 265.
- Shati AA. 2011. Effects of *Origanum majorana* L. on cadmium induced hepatotoxicity and nephrotoxicity in albino rats. *Saudi Med J.*, **32**(8): 797-805.
- Shi J, Yu J, Pohorly JE and Kakuda Y. 2003. Polyphenolics in grape seeds-biochemistry and functionality. *J Med Food.*, **6**(4): 291-9.
- Shinagawa FB, De Santana FC, Torres LRO and Mancini-Filho J. 2015. Grape seed oil: a potential functional food. *Food Sci Technol Campinas.*, **35**: 399-406.
- Shivananda NB, Dan Ramdath D, Marshall JR, Isitor G, Xue S and Shi J. 2011. Wound healing properties of the oils of *Vitis vinifera* and *Vaccinium macrocarpon*. *Phytother Res.*, **25**: 1201–1208.
- Singh RP, Murthy KN and Jayaprakasha GK. 2002. Studies on the antioxidant activity of pomegranate (*Punica granatum*) peel and seed extracts using *in vitro* models. *J Agric Food Chem.*, **50**: 81-86.
- Skipper A, Sims JN, Yedjou CG and Tchounwou PB. 2016. Cadmium chloride induces DNA damage and apoptosis of human liver carcinoma cells via oxidative stress. *Int J Environ Res Public Health*, **13**:1–10.
- Soliman MM, Attia HF, Hussein MM, Hassan ME and Ismail TA. 2013. Protective effect of N-acetylcysteine against titanium dioxide nanoparticles modulated immune responses in male albino rats. *Am J Immunol.*, **9**: 148-158.
- Soliman MM, Baiomy AA and Yassin MH. 2015. Molecular and histopathological study on the ameliorative effects of curcumin against lead acetate-induced hepatotoxicity and nephrotoxicity in Wistar rats. *Biol Trace Elem Res.*, **167**: 91-102.
- Takahashi O and Oishi S. 2003. Testicular toxicity of dietarily or parenterally administered bisphenol A in rats and mice. *Food Chem Toxicol.*, **41**: 1035-1044.
- Tchounwou PB, Yedjou CG, Patolla AK and Sutton DJ. 2012. Heavy metal toxicity and the environment. *Molecular, Clin Environ Toxicol.*, **3**: 133-164.
- Virk P, Elobeid M, Hamad S, Korany Z and Al-Amin M. 2013. Ameliorative effects of *Embllica officinalis* and *Rosmarinus officinalis* on cadmium-induced oxidative stress in Wistar rats. *J Med Plants Res.*, **7**: 805-818.
- Wang H, Chang M, Peng T, Yang Y, Li N, Luo T, Cheng Y, Zhou M, Zeng X and Zheng L. 2017. Exposure to cadmium impairs sperm functions by reducing catsper in mice. *Cell Physiol Biochem.*, **42**: 44-54.
- Yang XF, Ge YM, Jiang JQ, Xu ZY, Cui YH and Wang ZL. 2012a. Acute toxic effect of cadmium chloride in mice. *Chin J Vet Sci.*, **32**: 467-471.
- Yang XF, Ge YM, Zhang HT, Ning HM, Jiang JQ, Qi YH and Wang ZL. 2012b. Damaging effects of water-borne cadmium chloride on DNA of lung cells of immature mice. *Genet Mol Res.*, **11**: 4323-4329.
- Yildirim NC, Kandemir MF and Benzer F. 2011. Beneficial effects of grape seed extract against Cisplatin-induced testicular damage in rabbits. *Digest J Nanomat Biostruct.*, **6**: 155-159.