JJBS

Volume 14, Number 1, March 2021
ISSN 1995-6673
Pages 137 – 146
https://doi.org/10.54319/jjbs/140118

Jordan Journal of Biological Sciences

# Vegetative Morphological Variations within Some Egyptian *Amaranthus* L. Species

Wafaa K. Taia*, Azza A. Shehata, Manaser M. Ibrahim and Islam M. El-Shamy

[1]Faculty of Science, Botany and Microbiology Department, Alexandria University, Alexandria, Egypt

Received: November 8, 2019; Revised: January 14, 2020; Accepted: February 1, 2020

**Abstract**

Leaf and stem vegetative morphological characters have been studied in ten Egyptian *Amaranthus* species. Macro-and micro-morphological characters besides leaf venation and architecture have been studied. Twenty-seven main variable characters have been subjected as OTU's to numerical analysis. The resulted dendrograms (nearest neighbor, group average and farthest neighbor) divided the studied species into two categories with different subdivisions. From the most important morphological characters within the studied taxa are the places of red spot on the leaf surfaces and the type of leaf venation. These characters divided the studied taxa into two main groups: 1- the red spot; on the middle versus on the lower part of the leaf blades, 2- the leaf venation; reticulodromus versus brochidodromous.

**Keywords :** *Amaranthus*, Clustering analysis, Leaf characters, Macro-morphology, Micro-morphology, Stem characters, Venation.

## 1. Introduction

Genus *Amaranthus* L. is one of the widely distributed genera grown mainly in temperate and tropical climates, belonging to Amaranthaceae Juss., subfamily Amaranthoideae, tribe Amarantheae. Genus *Amaranthus* is the largest and has 70 annual or short lived perennial monoecious or dioecious species with worldwide distribution. It grows in roadsides, gardens and disturbed habitats with great morphological diversity between the species and even between individuals of the same species in response to environmental variables. *Amaranthus* has been faced with many opinions concerning species delimitation and identification since a long time ago. Thellung, (1914), Sauer, (1955), Aellen, (1961 and 1972), Cavaco, (1962) and Townsend, (1974 and 1985) have been faced with many difficulties in recognizing the different taxa while studying the morphological characters of the genus. Linnaeus, (1753) divided the genus into two groups; *Pentandri* and *Triandri* according to the number of stamens in the flower (which is the same as the number of perianth segments). This division of the genus has been accepted by Adanson, (1763) and Tournefort, (1794), while Dumortier, (1827) reclassified the *Amaranthus* species into two sections *Amaranthotypus* and *Blitopsis*, the former has monoecious plants with pentamerous flowers arranged in lax or dense spikes or panicles and circumscissile fruits. While section *Blitopsis* has dioecious plants with trimerous or pentamerous flowers arranged in axillary glomerules and has irregular ruptured indehiscent capsules.

Gordon, (1853) raised the inflorescence type to be a priori character in distinguishing the taxa under the genus and reclassified it into two subgenera; *Albersia* and *Euamaranthus*. And this reclassification has been accepted by both Kirschleger, (1857) and Bentham and Mueller, (1870), but they divided subgenus *Albersia* into three sections *Amblogyne*, *Euamaranthus*, and *Euxolus* according to the fruit whether bursts transversely or in an irregular manner or indehiscent. Afterward, Uline and Bray, (1894) classified the *Amaranthus* species into four groups; Amblogyne, Scleropus, Pyxidium, and Mega according to fruit characters. While Beck (1909) regrouped the *Amaranthus* species into three sections and Rouy, (1910) return back to the old classification of the species into two sections but renamed them *Euamaranthus* and *Pentrius*. Sauer, (1955), previously classified genus *Amaranthus* into two subgenera, differentiating only between monoecious and dioecious species, viz. *Amaranthus* and *Acnida* (L.) Aellen ex K. R. Robertson

The classification of *Amaranthus* is obscure due to the shortage of scrappy quantitative species-defining characteristics, besides the wide range of phenotypic flexibility, as well as introgression and hybridization involving weedy and crop species. This phenotypic variability led to nomenclature disorder and misapplication of names as seen in Mosyakin and Robertson, (1996), Costea *et al*., (2001) and Iamonico, (2009). Accordingly, the previous trials of classifications of the genus were faced with many difficulties in distinguishing and circumscription of the *Amaranthus* species. Mosyakin and Robertson, (1996) found that all the *Amaranthus* species must be divided into three subgenera; 1- *Acnida* (L.) Aellen ex K. R. Robertson with three sections; 2- *Albersia* (Kunth) Gren. and Godr. with four sections; and 3- *Amaranthus* with three sections and two subsections (Figure 1).

* Corresponding author e-mail: taia55taxonomy@hotmail.com.




**Figure 1.** Mosyakin and Robertson (1996) division of *Amaranthus* species

Till now the classification of the genus is still a matter of controversy and faced with great arguments. This work has been carried out to investigate the variations within the *Amaranthus* species grown in Egypt. According to Täckholm, (1974) the genus was represented by fourteen annual species; El Hadidi and El Hadidi, (1980) reported 10 species with 5 subspecies and 5 varieties, while according to Boulos, (1999, 2009) the genus was represented by eleven annual, rarely perennial species and five subspecies. These taxa growing, either as weeds in the cultivated areas or wildly in the roadsides. Ten of these species can be obtained from the field and subjected to analysis in this study to increase knowledge about the morphological variability between the species in the local region.

## 2. Materials and Methods

Ten *Amaranthus* species were collected from the wild population in Alexandria and Cairo during the years 2015 and 2016. The ten studied species arranged according to the system of infra-generic classification of Mosyakin and Robertson, (1996), their different collection sources are cited in tables 1.and 2 The collected specimens were identified using Täckholm, (1974) and Boulos, (1999). Taxonomic authorities for Latin names and synonymy of the species investigated in this study were based on either the Australian plant name index (APNI), Gray card index (GCI) or Index Kewensis (IK). Voucher specimens were kept in the Faculty of Science, Alexandria University Herbarium (ALEX) for vegetative morphological measurements and any further inspection.

**Table 1.** *Infra-generic* classification of *Amaranthus* species used in the present study

| Authors | Year | Characters | Ranks | Names | Further divisions |
|---|---|---|---|---|---|
| Linnaeus | 1753 | Perianth number | Groups | *Pentandri & Triandri* | -- |
| Adanson | 1763 | Perianth number | Groups | *Pentandri & Triandri* | -- |
| Tournefort | 1794 | Perianth number | Groups | *Pentandri & Triandri* | -- |
| Dumortier | 1827 | Sex of plant | sections | *Amaranthotypus & Blitopsis* | -- |
| Gorden | 1855 | Type of Inflorescence | Subgenera | *Albersia &Euamaranthus* | -- |
| Kirschleger | 1857 | Type of Inflorescence & Fruit characters | Subgenera | *Albersia (3 sections) & Euamaranthus* | *Amblogene, Euamaranthus & Euxalus* |
| Bentham | 1870 | Type of Inflorescence & Fruit characters | Subgenera | *Albersia (3 sections) &Euamaranthus* | *Amblogene, Euamaranthus & Euxalus* |
| Uline and Bray | 1894 | Fruit | Sections | *Amblogyne, Scleropus, Pyxidium & Megea* | -- |
| Beck | 1909 | Fruit | Sections | *Amblogene, Euamaranthus & Euxalus* | -- |
| Rouy | 1910 | Perianth parts | Sections | *Euamaranthus & Pentrius* | -- |
| Sauer | 1955 | Sex of plant | Subgenera | *Acnida & Amaranthus* | -- |
| Mosyakin and Robertson | 1996 | Sex of plant & Inflorescence | Subgenera | *Acnida, Amaranthus* & *Albersia* | *Acnida* = 3 sections<br>*Amaranthus* = 3 sections & 2 subsections<br>*Albersia* = 4 sections |

**Table 2.** Data of *Amaranthus* species collected in the present study and its taxonomic category after Mosyakin and Robertson, (1996)

| No. | Species | Collection | Locations | Subgenus | Sections | Sub- Sections |
|---|---|---|---|---|---|---|
| 1 | *A. albus* L. | 12 Jan. 2016,Wafaa Taia | Madinati,Cairo | *Albersia (Kunth) Gren. & Godr* | *Pyxdium oquin* in DC. | -- |
| 2 | *A. blitiodes* S. Watson | 11 Aug. 2015, Azza Shehata & Eslam El-Shamy | Fac. Agric., El-Shatby, Alexandria | *Albersia (Kunth) Gren. & Godr* | *Pyxdium oquin* in DC. | -- |
| 3 | *A. caudatus* L. | 11 Nov. 2015, Wafaa Taia | Madinet Nasr, Cairo. | *Amaranthus* | *Amaranthus* | *Amaranthus* |
| 4 | *A. graecizans* L. | 12 Dec. 2015, Wafaa Taia | . Smouha, Alexandria | *Albersia (Kunth) Gren. & Godr* | *Pyxdium Moquin* in DC. | -- |
| 5 | *A. hybridus* L. | 15 Nov. 2015, Azza Shehata & Eslam El-Shamy | Fac. Of Sci., El-Shatby, Alexandria | *Amaranthus* | *Amaranthus* | *Hybrida* |
| 6 | *A. lividus* L. | 1 Oct. 2015, Azza Shehata & Eslam El-Shamy | Fac. Of Sci., El-Shatby, Alexandria | *Albersia (Kunth) Gren. & Godr* | *Blitopsis Dumort* | -- |
| 7 | *A. retroflexus* L. | 14 Oct. 2015, Azza Shehata & Eslam El-Shamy | Fac. Of Sci., El-Shatby, Alexandria | *Amaranthus* | *Amaranthus* | *Amaranthus* |
| 8 | *A. spinosus* L. | 22 Oct. 2015, Azza Shehata & Eslam El-Shamy | Al-Agamy, Alexandria | *Amaranthus* | *Centrusa Griseb* | -- |
| 9 | *A. tricolor* L. | 10 Jun. 2015, Wafaa Taia | Al-Saha Square, Madinet Nasr, Cairo. | *Albersia (Kunth) Gren. & Godr* | *Pyxdium Moquin* in DC. | -- |
| 10 | *A. viridis* L. | 1 Oct. 2015, Manaser Ibrahim | Smouha, Alexandria | *Albersia (Kunth) Gren. & Godr* | *Blitopsis Dumort* | -- |

Stem color and habits were noticed in the field. At least ten individuals were used in this study, for morphological measurements and examinations. The fourth interned leaves from ten individuals in each species were subjected to measurements. For epidermal cell investigation using LM, fresh leaves were collected from each plant samples, painted with fingernail polish on both the abaxial and adaxial surfaces and allowed to dry. After drying, short clear cellophane tape was firmly pressed over the dried nail polish on the surfaces according to the method of Mbagwu *et al.,* (2008). Epidermal strips were taken from the median portion of matured leaves stained in alcoholic safranin and mounted in 50% glycerin jelly for microscopic examination. Photographs of good preparations were taken at a magnification of X400 objective using the light microscope (Olympus CX 31 with its built in camera).

For leaf venation, fresh leaves were immersed in 70% ethyl alcohol for a few days with several changes of alcohol to remove chlorophyll pigments. Leaf samples were washed by water. Leaves were carefully brushed out to obtain leaf skeleton, examined and photographed using Olympus stereomicroscope. For SEM, dried leaves were fixed directly to the copper stub with double-sided adhesive tape at abaxial and adaxial surfaces and coated for 5 minutes with gold in polaron JFC-1100E coating unit, then examined and photographed with JEOL JSM-5300 SEM at the EM unit, Faculty of Science, Alexandria University. The terminology of the leaf was according to Stace, (1984), and Barthlot *et al.,* (1998) to describe the epidermal cell features and leaf venation.

The characters used for clustering analysis were chosen according to be of a fixed taxonomic value and not affected by any environmental conditions. Consequently, 27 characters represented in Appendix (1) have been subjected to this analysis using Systat 12 program.

## 3. Results

The general external morphology of the studied species was illustrated in plate 1. The results were summarized in tables 3, 4 and 5, plates 1, 2, 3 and 4. *Amaranthus* species are annuals; except *A. hybridus*, *A. lividus* and *A. viridis* are perennials. The stem takes different shades of green and sometimes tinged with pink; erect, procumbent or prostrate. The stem has different length and width, slender, angular or flattened with moderate to dense branches and either glabrous or covered with hairs in various densities (Table 3). The leaves are green, pale green or tinged with red or reddish purple; spiral or alternate. The leaves are ovate to lanceolate, with the broader part in the middle, except in both *A. hybridus A. lividus* and *A. viridis* the wider part in the lower part of the blade. All the studied species have hairy leaf surfaces with different densities; except *A. lividus* the leaves are glabrous. The hairs are mostly multicellular uniseriate glandular (Plate 4f), but in *A. blitoides* the hairs are multicellular uniseriate tabular (Plate 3e) and in *A. tricolor* the hairs are multicellular uniseriate pointed (Plate 4g). Multicellular biseriate hairs noticed in *A. retroflexus* only beside the multicellular uniseriate ones (table 4). The leaf blade venation is either reticulodromous or brochidodromous (Plate 2a and b), with different primary veins thickness. The secondary veins varied from 7 to 15, irregular or regular and in few species, they are wider at the leaf blade apex. The marginal veins are complete or incomplete with incomplete aeriolar veins except in *A. albus*, the aeriolar veins are closed (table 4).

Leaf micro-morphological characters were summarized in table 5. The epidermal cells as shown by LM and SEM have elongated and irregular shapes in the abaxial surfaces, while they are polygonal and irregular in the adaxial surfaces (plates 2, 3, and 4).

The anticlinal walls are straight and elevated in all the species, except in *A. graecizans*, *A. hybridus* and *A.*

*tricolor* the anticlinal walls are undulated (Plate 3g - j and Plate 4g and h). The ornamentation of the periclinal walls is mostly striate, rugate, wrinkled or papillate with few or dense epicuticular secretions, except in *A. graecizans*, *A. hybridus* and *A. lividus.* While in *A. blitoides* the periclinal walls are smooth with few granulate epicuticular secretions (Plate 3c and d). The stomata are anomocytic or anisocytic (Plate 2 c, d, f and g), raised in all the studied species in both the abaxial and adaxial surfaces, except in *A. blitoides* it is sunken in the adaxial surface (Plate 3d). Meanwhile, the stomata are diacytic (Plate 2e and h) and superficial in both the abaxial and adaxial surfaces in *A. lividus* (Plate 4a and b).

The most valuable morphological variations are subjected to numerical investigations by taking 27 characters as OUT's in the ten studies taxa and the obtained dendrograms showed that, the three dendrograms split the studied taxa into two groups (GI and GII) at dissimilarity matrix 3 in the farthest neighbor linkage (F.N.); 5.3 in group average (G.A.) and 3.5 in the nearest neighbor (N.N.). In the nearest neighbor linkage (Figure 2a) and farthest neighbor linkage (Figure 2c) the dendrograms coincide in the division of the taxa; as both of them gather the taxa number 5, 9, 7, 6, 10 and 3 in one group and the taxa number 2, 8, 1 and 4 in the second group at different dissimilarity matrices as shown in table 6. In group average linkage method (Figure 2b) the two taxa (9 and 5) were separated in a one group (GII) and the rest of the taxa in a second group (GI). The three dendrograms gathered both taxa (3 and 10) together at a similarity index 1.012. The F.N. and G.A. gathered the taxa 9 and 5 at dissimilarity index 1.745 and 2.292; and the taxa 2 and 8 at dissimilarity index 3.623 and 1.675, respectively (Table 6).




**Figure 2a.** Nearest neighbor linkage **Figure 2b.** Group average linkage **Figure 2c.** Farthest neighbor linkage

**Figures 2a, b and c:** Dendrograms showing the three possible relationships among the ten studied *Amaranthus* species using Euclidean distance method and; **a.** Nearest neighbor linkage, **b.** Group average linkage and **c.** Farthest neighbor linkage methods. Cases studied: 1. *A. albus* ; 2. *A. blitiodes* ; 3. *A. caudatus* ; 4. *A. graecizans* ; 5. *A. hybridus* ; 6. *A. lividus* ; 7. *A. retroflexus* ; 8. *A. spinosus* ; 9. *A. tricolor* and 10. *A. viridis.*

**Table 3.** Stem morphological characters of the studied *Amaranthus* species

| No. | Species | Characters | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|
| | | Habit | Type | Length (cm) | Width (mm) | Shape | Color | Density of branches | Density of hairs |
| 1 | *A. albus* | Annual | Erect | 17-30 (23.5±5.44) | 0.2 – 0.4 (0.32±0.09) | Slender | Light green | Dense | Hairy |
| 2 | *A. blitiodes* Watson | Annual | Prostrate | 25-35 (28.6±3.78) | 0.1 - 0.2 (0.15±0.05) | Flattened | Greenish white | Moderate | Hairy |
| 3 | *A. caudatus* | Annual | Erect | 30-40 (36.6±2.07) | 0.2 - 0.5 (0.35±0.12) | Angular | Purple | Moderate | Hairy |
| 4 | *A. graecizans* | Annual | Procumbent | 14-30 (20.5±6.8) | 0.2 - 0.4 (0.3±0.08) | Angular | Red | Moderate | Sparsely hairy |
| 5 | *A. hybridus.* | Perennial | Erect | 40-55 (47.4±7.16) | 0.2 - 0.4 (0.32±0.09) | Angular | Green | Dense | Hairy |
| 6 | *A. lividus* | Perennial | Procumbent | 20-39 (34.8±5.67) | 0.1 - 0.3 (0.2±0.08) | Flattened | Green | Moderate | Hairy |
| 7 | *A. retroflexus* | Annual | Erect | 35-40 (36.8±1.92) | 0.2 - 0.4 (0.3±0.08) | Slender | Light Green | Moderate | Hairy |
| 8 | *A. spinosus* | Annual | Erect | 15-30 (25±7.07) | 0.3 - 0.6 (0.45±0.12) | Flattened | Green | Dense | Hairy |
| 9 | *A. tricolor* | Annual | Erect | 39-45 (41±2.54) | 0.2 - 0.3 (0.22±0.05) | Slender | Red | Moderate | Glabrous |
| | *A. viridis* | Perennial | Procumbent | 27-40 (36.2±6.37) | 0.2 - 0.4 (0.3±0.08) | Angular | Green or tinged purple | Moderate | Sparsely hairy |

± = Standard deviation values

**Table 4.** Leaf macro-morphological characters of the studied *Amaranthus* species

| No. | Species | Characters Arrangement | Color | Length (cm.) | Width (cm.) | Shape | Widest part | Spot | Surface | Hair type* |
|---|---|---|---|---|---|---|---|---|---|---|
| 1 | *A. albus* | Spiral | Pale Green | 0.9-3.1 (2.25±1.02) | 0.4-1.5 (0.97±0.45) | Lancolate | Middle | Middle | Sparsely hairy | MUG |
| 2 | *A. blitiodes* Watson | Alternate | Green with white spots | 1-2.4 (1.94±0.65) | 1-1.5 (1.32±0.16) | Ovate to Lancolate | Middle | Middle | Sparsely hairy | MUT |
| 3 | *A. caudatus* | Alternate | Green | 4-4.5 (4.20±0.20) | 1-2 (1.54±0.28) | Ovate to Lancolate | Middle | Middle | Hairy | MUG |
| 4 | *A. graecizans* | Alternate | Green tinged with red | 1.8-3.8 (2.77±1.02) | 1-1.9 (1.32±0.39) | Ovate to Lancolate | Middle | Middle | Sparsely hairy | MUG |
| 5 | *A. hybridus.* | Alternate | Green | 2-5.9 (4.52±1.66) | 1-4 (2.64±1.25) | Ovate | Lower | Lower | Wooly | MUG |
| 6 | *A. lividus* | Spiral | Green | 1.5-4.5 (2.70±1.29) | 1-3.5 (1.88±0.92) | Ovate | Lower | Lower | Glabrous | - |
| 7 | *A. retroflexus* | Alternate | Green tinged with red | 3-3.5 (3.18±0.21) | 1-1.5 (1.28±0.21) | Ovate to Lancolate | Middle | Middle | Hairy | MUG+ MBG |
| 8 | *A. spinosus* | Spiral | Green | 2.5-6 (4.32±1.47) | 2-3.8 (2.32±1.20) | Ovate to Lancolate | Middle | Middle | Hairy | MUG |
| 9 | *A. tricolor* | Alternate | Green or reddish purple | 3-4 (3.54±0.36) | 1-2 (1.54±0.36) | Ovate | Middle | Middle | Hairy | MUP |
| 10 | *A. viridis* | Alternate | Green tinged with purple | 2-5.4 (3.60±1.47) | 1-3.7 (2.58±0.94) | Ovate | Lower | Lower | Sparsely hairy | MUG |

*****Hair types: MUG = Muticellular Uniseriate Glandular, MUT = Muticellular Uniseriate Tubular, MBG = Muticellular biseriate Glandular and MUP = Muticellular Uniseriate Pointed, ± = Standard deviation values.

**Table 4 (Cont.)** Leaf macro-morphological characters of the studied *Amaranthus* species

| No. | Species | Characters | | | | | |
|---|---|---|---|---|---|---|---|
| | | Types of venation | Size of 1$^{ry}$ veins | Type of 2$^{ry}$ veins | | Marginal veins | Areolar Veins |
| | | | | Spaces | No. | | |
| 1 | *A. albus* | Reticulodromous | Moderate | Wide at apex | 8 | Incomplete | Closed |
| 2 | *A. blitiodes* Watson | Brochidodromous | Thin | Wide at apex | 12 | Incomplete | Incomplete |
| 3 | *A. caudatus* | Reticulodromous | Thick | Irregular | 15 | Incomplete | Incomplete |
| 4 | *A. graecizans* | Brochidodromous | Thick | Wide at base | 7 | Complete | Incomplete |
| 5 | *A. hybridus* | Brochidodromous | Thick | Wide at apex | 11 | Complete | Incomplete |
| 6 | *A. lividus* | Brochidodromous | Thick | Irregular | 11 | Incomplete | Incomplete |
| 7 | *A. retroflexus.* | Brochidodromous | Thick | Wide at apex | 9 | Complete | Incomplete |
| 8 | *A. spinosus* | Brochidodromous | Moderate | Irregular | 11 | Complete | Incomplete |
| 9 | *A. tricolor* | Reticulodromous | Thick | Irregular | 8 | Complete | Incomplete |
| 10 | *A. viridis* | Brochidodromous | Thick | Regular | 14 | Incomplete | Incomplete |



a) A . albus b) A . blitoides c) A . caudatus d) A . graecizans

e) A . hybridus f) A . lividus g) A . retroflexus h) A . spinosus

i) A . tricolor j) A . viridis

**Plate 1.** (a-j) Photo-micrographs showing the general external morphological features of the ten studied *Amaranthus* species; a) *A. albus*, b) *A. blitoides*, c) *A. caudatus*, d) *A. graecizans*, e) *A. hybridus*, f) *A. lividus*, g) *A. retroflexus*, h) *A. spinosus*, i) *A. tricolor*and and j) *A. viridis*, respectively.




a) Reticulodromous venation b) Brochidodromous venation




Anomocytic stomatal type Anisocytic stomatal type Diacytic stomatal type

**Plate 2.** (a and b) Photo-micrographs showing the types of the leaf venation, (c-e) Photo-micrographs, (f-h) drawings showing the types of the leaf stomatal complex for the studied *Amaranthus* species**.**

**Table 5.** Leaf micro-morphological characters of the studied *Amaranthus* species

| No. | Species | Characters | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|
| | | Shape of epidermal cell | | Epidermal wall pattern | | | Stomata type | | Stomata level | |
| | | Abaxial | Adaxial | Anticlinal wall | Periclinal wall | Epicuticuler secretion | Abaxial | Adaxial | Abaxial | Adaxial |
| 1 | *A. albus* | Elongated/ Irregular | Polygonal/ Irregular | Straight/ Elevated | Striate | Few granules | Anomo- Aniso- cytic | Anomo- Aniso-cytic | Raised | Raised |
| 2 | *A. blitiodes* Watson | Irregular | Polygonal/ Irregular | Straight/ Elevated | Smooth | Few granules | Anomo- Aniso- cytic | Anomo- Aniso-cytic | Raised | Sunken |
| 3 | *A. caudatus* | Sinuate/ Irregular | Polygonal/ Irregular | Straight/ Elevated | Rugate | Few granules | Anomo- Aniso- cytic | Anomo- Aniso-cytic | Raised | Raised |
| 4 | *A. graecizans* | Sinuate/ Irregular | Irregular | Undulate/ Elevated | Rugate | Absent | Anomo-cytic | Anomo-cytic | Raised | Raised |
| 5 | *A. hybridus* | Sinuate Irregular | Polygonal/ Irregular | Undulate/ Elevated | wrinkled | Absent | Anomo- Aniso- cytic | Anomo- Aniso- cytic | Raised | Raised |
| 6 | *A. lividus* | Elongated/ Irregular | Regular | Straight/ Elevated | Striate | Absent | Anomo- Aniso-Dia- cytic | Anomo- Aniso-Dia- cytic | Superficial | Superficial |
| 7 | *A. retroflexus* | Elongated | Isodiametric/ Irregular | Undulate/ Elevated | wrinkled | Dense granules | Anomo- Aniso-cytic | Anomo- Aniso- cytic | Raised | Raised |
| 8 | *A. spinosus* | Elongated | Polygonal/ Irregular | Undulate/ Elevated | Papillate | Dense granules | Anomo- Aniso-cytic | Anomo- Aniso- cytic | Raised | Raised |
| 9 | *A. tricolor* | Elongated/ Irregular | Elongated/ Irregular | Undulate/ Elevated | Striate | Dense granules | Anomo- Aniso-cytic | Anomo- Aniso- cytic | Raised | Raised |
| 10 | *A. viridis* | Elongated/ Irregular | Elongated/ Irregular | Straight/ Elevated | Striate | Few granules | Anomo- Aniso- cytic | Anomo- Aniso- cytic | Raised | Raised |

**Table 6.** Joining Euclidean distances gathered for the ten studied *Amaranthus* species according to the linkage methods of data analysis using Systat 12 program

| Nearest neighbor linkage method | | | | Group average linkage method | | | | Farthest neighbor linkage method | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| Clusters Joining | | Distance | No. of Members | Clusters Joining | | Distance | No. of Members | Clusters Joining | | Distance | No. of Members |
| Case 10 | Case 3 | 1.012 | 2 | Case 10 | Case 3 | 1.012 | 2 | Case 10 | Case 3 | 1.012 | 2 |
| Case 6 | Case 10 | 1.043 | 3 | Case 7 | Case 6 | 1.049 | 2 | Case 7 | Case 6 | 1.049 | 2 |
| Case 7 | Case 6 | 1.049 | 4 | Case 8 | Case 1 | 1.081 | 2 | Case 8 | Case 1 | 1.081 | 2 |
| Case 8 | Case 1 | 1.081 | 2 | Case 8 | Case 4 | 1.302 | 3 | Case 8 | Case 4 | 1.437 | 3 |
| Case 8 | Case 4 | 1.167 | 3 | Case 10 | Case 7 | 1.313 | 4 | Case 10 | Case 7 | 1.632 | 4 |
| Case 2 | Case 8 | 1.297 | 4 | Case 9 | Case 10 | 1.653 | 5 | Case 9 | Case 5 | 1.745 | 2 |
| Case 7 | Case 9 | 1.357 | 5 | Case 2 | Case 8 | 1.675 | 4 | Case 2 | Case 10 | 2.011 | 5 |
| Case 7 | Case 2 | 1.432 | 9 | Case 9 | Case 5 | 2.292 | 6 | Case 8 | Case 2 | 3.623 | 8 |
| Case 5 | Case 7 | 1.745 | 10 | Case 9 | Case 2 | 3.088 | 10 | Case 8 | Case 9 | 5.422 | 10 |

**Cases studied:** 1. A . albus; 2. A . blitiodes; 3. A . caudatus; 4. A . graecizans; 5. A . hybridus; 6. A . lividus; 7. A . retroflexus; 8. A . spinosus; 9. A . tricolor and 10. A .viridis .




**Plate 3**. (a-j) SEM photo-micrographs showing types of stomata and shape of epidermal cells on both abaxial and adaxial surfaces of the leaf for *A. albus, A. blitoides, A. caudatus, A. graecizans,* and *A. hybridus***,** respectively.




**Plate 4.** (a-j) SEM photo-micrographs showing types of stomata and shape of epidermal cells on both abaxial and adaxial surfaces of the leaf for *A. lividus, A. retroflexus, A. spinosus, A. tricolor* and *A. viridis*, respectively.

## 4. Discussion

*Amaranthus* is a cosmopolitan genus with annual monoecious or dioecious species. Species within this genus are under taxonomic investigations due to their high phenotypic variability (Mosyakin and Robertson, 2003; Iamonico, 2009 and 2015). The first comprehensive taxonomical division of the genus was that of Dumortier, (1827) who divided its species into two sections. Classification of the genus has been faced with many opinions and divisions afterward; from the most reasonable division is that of Mosyakin and Robertson, (1996). Accordingly, the vegetative morphological variation within the Egyptian *Amaranthus* has been investigated in this work. Davis and Heywood, (1973) pointed out that any stable character can be used in systematic. Subsequently, the most stable characters have been subjected to numerical analysis to investigate the relationship between the studied species.

According to Dumortier, (1827), the studied taxa were classified under the two sections; *Amaranthotypus* and *Blitopsis*; as follows: *A. caudus, A. hybridus, A. retroflexus,* and *A. spinosus* under section *Amaranthotypus*. While section *Blitopsis* includes *A. albus, A. blitoides, A. graecizans, A. lividus, A. tricolor* and *A. viridis*. The results found in the present study did not coincide with Dumortier's classification of the species. The obtained dendrograms divided the studied taxa into two groups in the three clustering methods (N.N., G.A. and F.N.). The three dendrograms put *A. albus* beside *A. spinosus* (1 and 8) at dissimilarity index 1.081, as well as *A. caudatus* beside *A. viridis* (3 and 10) at dissimilarity matrix 1.012. The species *A. lividus* and *A. retroflexus* (6 and 7) are closely met at dissimilarity matrix 1.049 in the three dendrograms. These gatherings of the species are logic due to the morphological similarities between every two species, in spite of gathering species between two different Dumortier's sections. This result may be in acceptance of Mosyakin and Robertson, (1996) who divided all the *Amaranthus* species into three subgenera; 1- *Acnida* (L.) Aellen ex K. R. Robertson with three sections; 2- *Albersia* (Kunth) Gren. and Godr. with four sections and 3- *Amaranthus* with three sections and two subsections.

As *Acnida* taxa were not represented in the flora of Egypt, accordingly the three resulted dendrograms revealed that the studied taxa can be divided into two subgenera, with many sections and *Acnida* is excluded. We can explain this division as it was according to the vegetative morphological characters only. From the most important morphological characters within the studied taxa are the places of red spot on the leaf surfaces and leaf venation, which divide the studied taxa into two groups, red spot on the leaf surfaces on the middle versus on the lower part of the leaf blades besides the leaf venation, reticulodromus versus brochidodromous. The species under these groups did not coincide with those of Dumortier, (1827). In spite of that, the results obtained were in partial acceptance with that of Waseltov *et al.*, (2018) who found that both *A. caudatus* and *A. hyvridus* were closed to each other in the phylogenetic tree, as well as *A. albus* and *A. blitoides*. Most of the taxonomical works on this genus were classified according to the floral characters and sex of the plants (Mosyakin and Robertson, 2003 and Iamonico, 2009 and 2015). As a conclusion the studied *Amaranthus* species can be divided into two main

groups based on their vegetative morphological characters only.

## References


Adanson M. 1763. **Familles des Plantes.**, Vincent, Paris.

Aellen P. 1961, **Die Amaranthaceae Mitteleuropas**. Carl Hanser Verlag, München.

Aellen P. 1972. In: Rechinger K. (ed.), Amaranthaceae. **Flora Iranica 91**, Graz-Austria.

Barthlott W; Neinhuis C; Cutler D; Ditsch F; Meusel I; Theisen I and Wilhelmi H. 1998. Classification and terminology of plant epicuticular waxes. *Bot J Linn Soc*, **126**:237-260.

Beck GE. 1909**. Amaranthaceae**. In: Reichenbach, L.**,** and Reichenbach, HG (ed.), Icones Florae Germanicae et Helveticae. Lipsiae et Gerae pp. 91-174.

Bentham G. and Mueller F. 1870. **Flora Australiensis** vol.5. L. Reeve and co., London.

Boulos L. 1999. **Flora of Egypt**. Vol. 1. Azollaceae-Oxalidaceae, Al Hadara Publishing, Cairo, Egypt.

Boulos, L. 2009. Flora of Egypt. Checklist Al-Hadara Publishing, Cairo.

Cavaco A. 1962. **Les Amaranthaceae de l'Afrique au sud du tropique du Cancer et de Madagascar**. Mémoires du Muséum National d'Histoire Naturelle, sér. B, *Botanique* **13**: 1-254.

Costea M; Sanders A and Waines G. 2001. Notes on some little known *Amaranthus* taxa (Amaranthaceae) in the United States. *Sida*, **19**: 931-974.

Davis PH and Heywood VH. 1973. **Principles of Angiosperm Taxonomy**. Krieger, New York.

Dumortier, BCJB. 1827. **Florula Belgica Operis Majoris Prodromus**, Belgique.

El Hadidy, MN and El Hadidy AMH. 1980. Family 57. Amaranthaceae in M N. El Hadidi (Ed.). Flora of Egypt. *Taeckholmia add ser*.**1**:13-92.

Gordon D.1853. **Florula Juvenalis III** *Mém Acad Sci et Lettr de Montp*, .

Iamonico D. 2009. Invasive status and presence of *Amaranthus polygonoides* L. (Amaranthaceae) in Italy, with notes on its taxonomy and morphology. *Fl Medit.*, **19**: 233-239.

Iamonico D. 2015. Taxonomic revision of the genus *Amaranthus* (Amaranthaceae) in Italy. *Phytotaxa*, 199(1): 1-84.

Kirscleger F. 1857. **Flore d'Alsace et des contrées,** limitrophes Vol. 2, Les monochlamydées, Les monocotylées, Les cryptogames vasculaires.

Linneaus C. 1753. **Species Plantarum**. Vol. 2, Holmiae.

Mbagwu FN; Nwachukwu CU and Okoro OO. 2008. Comparative Leaf Epidermal Studies on *Solanum macrocarpon* and *Solanum nigrum*. *Res J Bot.*, **3**(1): 45-48.

Moquin-Tandon, CHBA (1849) **Amaranthaceae Juss**. In: De Candolle A (Ed.). Prodromus Systematis Regni Vegetabilis, Sumptibus Victois Masson, Parisiis, **13**(1): 231-424.

Mosyakin SL and Robertson KR. 1996. New infrageneric taxa and combinations in *Amaranthus* (Amaranthaceae). *Ann Bot Fenn*, **33**:275-281.

Mosyakin SL and Robertson KR. 2003. *Amaranthus*. In: **Flora of North America North of Mexico**, Vol. **4**, Oxford University Press.

Rouy G. 1910. **Flora of France ou description des plante vol. XII**, Paris.

Sauer JD. 1955. Revision of the dioecious *Amaranths*. *Madroño*, **13**: 5-46.

Stace CA. 1984. The taxonomic importance of the leaf surface. In: Heywood VH and Moore DM (eds.) **Current concepts in plant taxonomy**, Academic Press, London, U.K.

Täckholm V. 1974. **Students' Flora of Egypt**, second ed., Cairo University

Thellung A. 1914. *Amaranthus*. In: Ascherson P and Graebner P (eds.), **Synopsis der Mitteleuropaischen Flora 5**, Leipzg.

Tournefort JP. 1794. **Institutiones rei Herbariae**. Editio Altera, Paris.

Townsend CC. 1974. Amaranthaceae. In: Nasir E and Ali SI (Eds.) **Flora of West Pakistan** 7. Ferozsons Press, Rawalpindi

Townsend CC. 1985. Amaranthaceae **In: Polhill RM (ed.). Flora of Tropical East Africa**. Published on behalf of the East African Government.

Uline EB and Bray WL. 1894. A preliminary synopsis of the North American species of *Amaranthus*. *Bot Gaz.*, **19**: 313-320.

Waselkov KE; Boleda A and Olsen KM. 2018. A Phylogeny of the Genus *Amaranthus* (Amaranthaceae) Based on Several Low-Copy Nuclear Loci and Chloroplast Regions. *Syst Bot.*, **43**(2): 439-458.

**Appendix 1.** Types of characters (OTU's) used in the numerical analysis and its ranking

| No. | | Characters | Ranking of Multistated Qualitative | | | | | |
|---|---|---|---|---|---|---|---|---|
| | | | 1 | 2 | 3 | 4 | 5 | 6 |
| 1 | | Habit | Annual | Perennial | | | | |
| 2 | | Type | Erect | Procumbent | Prostrate | | | |
| 3 | Stem characters | Length (cm) | Multistated Quantitative | | | | | |
| 4 | | Width (mm) | | | | | | |
| 5 | | Shape | Flattened | Angular | Slender | | | |
| 6 | | Color | Light Green | Greenish white | Green | Green tinged with purple | Purple | Red |
| 7 | | Density of branches | Moderate | Dense | | | | |
| 8 | | Density of hairs | Glabrous | Sparsely hairy | Hairy | | | |
| 9 | | Arrangement | Alternate | Spiral | | | | |
| 10 | | Color | Pale green | Green | Green with white spots | Green tinged with red | Green tinged with purple | Green or reddish purple |
| 11 | Leaf macro-characters | Length (cm) | Multistated Quantitative | | | | | |
| 12 | | Width (cm) | | | | | | |
| 13 | | Shape | Ovate | Ovate to Lancolate | Lancolate | | | |
| 14 | | Widest part | Middle | Lower | | | | |
| 15 | | Spot | Middle | Lower | | | | |
| 16 | | Surface | Glabrous | Sparsely hairy | Hairy | Wooly | | |
| 17 | | Hair type* | MUP | MUT | MUG | MUG+MBG | | |
| 18 | | Types of venation | Reticulodromous | Brochidodromous | | | | |
| 19 | | Size of $1^{ry}$ veins | Thin | Moderate | Thick | | | |
| 20 | | Type of $2^{ry}$ veins space | Regular | Wide at apex | Wide at base | Irregular | | |
| 21 | | Number of $2^{ry}$ veins | Multistated Quantitative | | | | | |
| 22 | Leaf micro-characters | Marginal veins | Complete | Incomplete | | | | |
| 23 | | Areolar veins | Closed | Incomplete | | | | |
| 24 | | Abaxial epidermal cells | Irregular | Elongated/Irregular | Sinuate/Irregular | | | |
| 25 | | Adaxial epidermal cells | Irregular | Elongated/Irregular | Polygonal/Irregular | Isodiametric/Irregular | Polygonal/regular | |
| 26 | | Abaxial stomatal type | Anomo-cytic | Anomo-Aniso-cytic | Anomo-Aniso- Dia-cytic | | | |
| 27 | | Marginal veins | Complete | Incomplete | | | | |