

# The one-pot synthesis of some bioactive pyranopyrazoles and evaluation of their protective behavior against extracellular H<sub>2</sub>O<sub>2</sub> and SNP in *T. Thermophila*.

Boutaina Addoum<sup>1</sup>, Bouchra El khalfi<sup>1</sup>, Reda Derdak<sup>1</sup>, Souraya Sakoui<sup>1</sup>, Abdelhakim Elmakssoudi<sup>2</sup>, and Abdelaziz Soukri<sup>1,\*</sup>

<sup>1</sup>Laboratory of Physiopathology, Genetics Molecular and Biotechnology (PGMB), Department of Biology, Faculty of Sciences Ain Chock, Research Center of Health and Biotechnology, Faculty of Sciences Ain Chock, Hassan II University, B.P 5366 Maarif, Casablanca, Morocco, <sup>2</sup>Laboratory of Organic Synthesis, Extraction, and Valorization (OSEV), Department of Chemistry, Faculty of Sciences Ain Chock, Hassan II University, B.P 5366 Maarif, Casablanca, Morocco.

Received Feb 29, 2020; Revised June 13, 2020; Accepted June 26, 2020

## Abstract

This study aims to evaluate the antioxidant activity of pyranopyrazoles derivatives synthesized *via* multicomponent reaction and characterized by spectroscopic techniques (<sup>1</sup>H-NMR, <sup>13</sup>C-NMR and IR). All the synthesized compounds were screened for their anti-stress properties *in vivo* by using as a test organism the eukaryotic cells "*Tetrahymena*" exposed to oxidative and nitrosative stress. Thereafter the antioxidant activity *in vitro* is performed by using the 2,2-diphenyl-1-picrylhydrazyl (DPPH) scavenging assays.

Our results clearly showed the protective behavior of the compounds studied (**5a** and **5b**), especially against H<sub>2</sub>O<sub>2</sub>-induced stress. On the other hand, the *in vitro* bioassays reveal that the **5a** derivative possesses significant antioxidant activity much greater than ascorbic acid (Vitamin C). All these findings confirmed the prominent antioxidant behavior of the synthesized compounds **5a** and **5b**. However, further studies are needed to describe the mechanism action and the effect of substitution on the activity of these compounds.

**Keywords:** Pyranopyrazoles, *Tetrahymena*, Stress, Antioxidant Activity, Synthesis.

## 1. Introduction

Oxidative stress is defined as the aggression of cellular constituents associated with an unbalance between reactive species and the antioxidant system (Kumar *et al.*, 2015). The cells are damaged when they are emerged by ROS and RNS, it includes nucleic acids (DNA, RNA), proteins and lipids sequestration, which play an important role in the pathogenesis of several diseases (Fourrat *et al.*, 2007).

Recently, a large variety of synthetic products were aberrantly exploited as efficient antioxidant drugs, which supports the inner antioxidant defense system (SOD, GLx, Glutathione, Vitamin C) and overcomes many free radical-related diseases including cancer, diabetes, kidney failure, Alzheimer's, and Parkinson's (Pisoschi and Pop, 2015).

Among these efficient and available synthetic products, we found the pyranopyrazoles derivatives (Dove P, 2016) that are considered as primeval measuring sticks for expansive preclinical and pharmacological investigations (Debasis, Banerjee, and Mitra, 2014) due to their wide-ranging activities such as antioxidant, antibacterial, antiviral, antifungal, anti molluscidal and analgesic behavior (Mamaghani and Hossein Nia, 2019).

This study began with the synthesis of pyranopyrazoles derivatives *via* a multicomponent reaction (MCR)

approach (Zahouily *et al.*, 2005) starting from aromatic aldehyde, ethylacetoacetate, malononitrile and hydrazine hydrate in the presence of Na<sub>2</sub>CaP<sub>2</sub>O<sub>7</sub> (Tekale *et al.*, 2013) as a catalyst in a green solvent (water) (Maleki *et al.*, 2016). Then the bioactivity of these compounds was screened *in vivo* and *in vitro*. In this regard, the well-studied protozoan *Tetrahymena thermophila* (Mar *et al.*, 2017; Sugimoto *et al.*, 1960) was subjected to oxidative and nitrosative stress then was treated with these compounds. This protozoan can be easily cultured and maintained under laboratory conditions (Doerder and Brunk, 2012) and their sensitivity to stress (Zhang *et al.*, 2015) exposure encouraged several researchers to consider this freshwater ciliate as a standard test organism to understand many cellular and molecular processes (Huang *et al.*, 2016). On the other side, these compounds were subjected also to the DPPH test to confirm their protective potential *in vitro* compared to the ascorbic acid as a potent antioxidant.

To be specific, the protective effects of pyranopyrazoles on *T. thermophila* have not been evaluated so far. Hence, we tested, examined and discussed for the first time the effects of these compounds on the physiological and kinetic parameters of *T. thermophila* cells exposed to oxidative stress, however, further studies are needed such as the evaluation of their

\* Corresponding author e-mail: ab.soukri@gmail.com.

effect on the specific activity of some antioxidant enzymes like: catalase (CAT), superoxide dismutase (SOD) and GAPDH (Glyceraldehyde 3-phosphate dehydrogenase).

## 2. Experimental


### 2.1. Material and methods

#### 2.1.1. Instruments

All the chemicals were purchased from Sigma Aldrich and used without any further purification. The  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra were recorded on a Bruker 300 MHz in  $\text{DMSO-d}_6$  as a solvent. Chemical shifts are given in parts per million ( $\delta$ -scale). The X-ray diffractometer (XRD) pattern of the diphosphate (DIPH)  $\text{Na}_2\text{CaP}_2\text{O}_7$  was recorded on a Bruker D8 Advance X-ray diffractometer with  $\text{Cu-K}\alpha$  radiation. The samples were analyzed by FT-IR spectroscopy (using a SCHIMADZU IRAffinity-1S in the range of  $4000\text{--}400\text{cm}^{-1}$ ). Analytical thin-layer chromatography was performed with Silica on TLC Alu foils. Visualization of the developed chromatogram was performed by UV light (254nm). The melting point of the pyranopyrazoles derivatives was determined using a Buchi 510 apparatus.

#### 2.2. Procedure for synthesis of pyranopyrazoles derivatives **5a** and **5b**.

The catalyst  $\text{Na}_2\text{CaP}_2\text{O}_7$  (20mol %) was added to a mixture of the aromatic aldehyde **1** (salicylaldehyde **1a**, *p*-anisaldehyde **1b**) (1 mmol), ethylacetoacetate **2** (1mmol), malononitrile **3** (1.2 mmol), hydrazine hydrate **4** (2 mmol), and 1 ml of water in a flask fitted with a reflux condenser. The resulting mixture was heated to reflux (an oil bath) with stirring for 20 min. Acetone (2 ml) was added and the mixture stirred for 2 min. The catalyst was removed by filtration and then the resulting crude reaction mixture was poured onto crushed ice and precipitated solid was collected and recrystallized using 96% ethanol to afford pyranopyrazoles **5a** and **5b** (Maleki *et al.*, 2016). The products were well characterized by their melting points  $^1\text{H}$ ,  $^{13}\text{C}$  NMR, and IR spectroscopy.



**Figure 1.** The synthesis reaction catalyzed by the  $\text{Na}_2\text{CaP}_2\text{O}_7$ .

### 2.3. Biological Evaluation: microorganisms' growth and toxicological bioassays

#### 2.3.1. Reagents


Peroxide hydrogen and SNP were purchased from Himedia. The culture media such as yeast extract and Tryptone were obtained from Biokar, DPPH, and ascorbic acid were obtained from Sigma-Aldrich. The pyranopyrazoles were synthesized in the Laboratory of Organic Synthesis, Extraction and Valorization according to the references and their structures are summarized in figure 1.

#### 2.3.2. Strains and growth conditions

Before the experiments, the strains *Tetrahymena Thermophila* (SB 1969) were grown axenically in the stock medium of PPYE (1.5% of protease peptone and 0.25% of yeast extract, autoclaved in high-pressure at  $120^\circ\text{C}$  for 30 min before use. The culture medium is inoculated with 1 % (v / v) of *Tetrahymena Thermophila* then incubated at  $32^\circ\text{C}$  (Addoum *et al.*, 2018; Mar, El Khalfi, and Soukri, 2018).

#### 2.3.3. Cells exposed to stress

To evaluate the effect of stress on protozoan growth, the cultures of *T. thermophila* are treated respectively with 0.4 mM of  $\text{H}_2\text{O}_2$  and 1.8mM of SNP after 24h of incubation as reported previously in our previous work (Addoum *et al.*, 2018). Each sample (treated and control) was maintained at  $32^\circ\text{C}$  for 72h and accompanied by an optical microscope display (A. KRÜSS Optronic) then a cell counting was performed to evaluate the time and number of generation (figure 2).



**Figure 2.** General protocol for stress induction. Cells were grown at  $32^\circ\text{C}$  in PPYE medium containing  $\text{H}_2\text{O}_2$  reagent (0.3mM) or SNP reagent (1.8mM).

#### 2.3.4. Growth measurement

The growth kinetics of *Tetrahymena* was established by daily monitoring of the optical density OD at a wavelength of 600 nm during 120h.

#### 2.3.5. Morphological changes

To observe the morphology of *T. Thermophila* in different experimental conditions (treated and untreated), fresh samples were removed, examined and images were taken using the A. KRÜSS Optronic microscope (magnifications x10).

#### 2.3.6. Cells densities

To determine the cell viability an aliquot of 1ml was taken immediately from the treated cultures and the control then supplemented with a suitable volume of formaldehyde (2-4%). After fixation, the relative cell number was determined by using a hemocytometer counting chamber (Malassez double cells).

#### 2.3.7. Growth Curves

Growth and cell viability curves of *T. Thermophila* were produced based on time-OD and cell density-time. To produce growth curves, flasks of growth medium were inoculated with *Tetrahymena* cells and followed different treatment (stress, antioxidant treatment, and control). During the culture period, growth (OD) and cell number were measured every 24 hours up to 120h. Three biological replicates were included in this study. The obtained values were used to produce the *T. Thermophila* curves and analyzed via the GraphPad prism 8:0.

### 2.4. Chemicals

#### 2.4.1. Stock solution

A stock solution of pyranopyrazoles (**10mg/mL**) was prepared in dimethylsulfoxide (**DMSO**) (0.01%) (Reddy et al., 2019). Then we prepared a gamut of dilutions from 10-0.6 mg/ML. The prepared molecules were stored at ambient temperature in obscurity.

#### 2.4.2. Toxicity assays

For each derivative two independent tests were performed; we started with the turbidity test that consists to add 5µl of each dilution separately to the culture medium, then we inoculate 1% (v/v) a calibrated inoculum ( $1.5 \times 10^5$  cells/mL) of protozoan *Tetrahymena*. Under this experimental condition, the protozoan is maintained at 32°C for 72h. Ultimately, we retain the minimum inhibitory concentration (MIC) to evaluate their protective effects against stress conditions. This concentration was referred to as the lowest concentration of the tested compound that yields a visible growth of the protozoan on the liquid medium tubes. On the other side, this test is confirmed by the investigation of total cells number via microscopic visualization (X10). Estimates of cell viability were reported as a percentage of cell numbers (individuals/ml).

#### 2.5. Antioxidant treatment

To assess the effect of pyranopyrazoles on ROS production; the appropriate stock solution and the test dilutions were selected. Five nominal test concentrations were chosen (table1) and added to a ciliate-containing

flask (100 ml of PPYE medium). The treatment with pyranopyrazoles was done at the beginning (t = 0) before the transplantation of *Tetrahymena* cells (105cell/ml). All experiments were performed with protozoan cells in the exponential growth phase. The optical densities of the protozoan are measured by a UV-visible spectrophotometer (Jenway 7315 UV) as described in the literature (Addoum et al., 2018; Errafiy, Ammar, and Soukri, 2013).

Simultaneously the determination of absorbance was accompanied by microscopic imaging analysis and we calculated also the cell densities via hemocytometer technique as described previously in this paper.

In each bioassay, we used two controls: the negative control (Zero mg/mL concentration of pyranopyrazoles), and the positive control (living cells not exposed to stress but treated with pyranopyrazoles). Each concentration was conducted in triplicate.

#### 2.6. DPPH scavenging activity

The detection of the anti-radical activity of pyranopyrazoles was performed by the method of DPPH (El-borai et al., 2016). Various concentrations of selected compounds were mixed with about 1ml of methanol solution then placed in a tube which was added 4ml of DPPH prepared in methanolic solution (0.004%). Then, the tubes are maintained at obscurity for 30 min. The absorbance was screened at 517 nm via a UV spectrophotometer; this reduction was accompanied by the color change from purple to yellow. We used as standard various concentrations of the ascorbic acid prepared in distilled water (El-borai et al., 2016).

The percentage inhibition of free radical DPPH (I%) was calculated by the following formula:

$$I\% = (A_{\text{control}} - A_{\text{sample}} / A_{\text{control}}) \times 100$$

Where  $A_{\text{control}}$  reflected the absorbance of the ascorbic acid and  $A_{\text{sample}}$  is the absorbance of pyranopyrazoles products.

#### 2.7. Statistical Analysis

The statistical data are expressed as mean +/- SD for three successive measurements by using Graph Pad Prism software 8.0. For statistical significance, we used Student's t-test and the P-value was fixed in  $\leq 0.05$  for a statistically significant difference.

### 3. Results

#### 3.1. Chemistry

##### 3.1.1. Synthesis

The current study described the catalyzed synthesis of some biologically active pyranopyrazoles by using an ecofriendly catalyst, and these compounds are prepared via a one-pot reaction depicted in scheme 2. Primarily, we endeavored to the catalyst four-component commercially available to condense *in situ*; aromatic aldehyde **1**, ethylacetoacetate **2**, hydrazine hydrate **3**, and malononitrile **4** in the water at reflux temperature. The pyranopyrazoles **5a** and **5b** were obtained in high yields (87% and 98 %, respectively).

The structures of the synthesized compounds were confirmed by using spectroscopic techniques.





### 3.1.2. Characterization Spectroscopic data of synthesized products 5a and 5b

**6-Amino-3-methyl-4-(2-hydroxyphenyl)-1,4-dihydropyran[2,3-*c*] pyrazole-5-carbonitrile 5a:** White solid; m.p. 215-218 °C; <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>, ppm): δ 1.9 (s, 3H), 4.56 (s, 1H), 6.70 (2H), 6.94-7.18 (m, 5H), 11.00 (s, 1H); <sup>13</sup>C NMR (75 MHz, DMSO-*d*<sub>6</sub>, ppm): δ 9.85, 28.64, 55.06, 104.95, 115.1, 120.79, 123.53, 124.25, 127.55, 128.93, 136.52, 148.39, 159.09, 160.08; IR (KBr, cm<sup>-1</sup>): 3448, 3419, 3352, 2189, 1660, 1610 (Kanagaraj and Pitchumani, 2010).

**6-Amino-3-methyl-4-(4-methoxyphenyl)-1,4-dihydropyran[2,3-*c*] pyrazole-5-carbonitrile 5b:** White solid; m.p. 210-212 °C; <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>, ppm): δ 1.78 (s, 3H), 3.72 (s, 3H), 4.54 (s, 1H), 6.83 (2H), 6.86-7.08 (m, 4H), 12.09 (s, 1H); <sup>13</sup>C NMR (75 MHz, DMSO-*d*<sub>6</sub>, ppm): δ 9.66, 35.51, 55.00, 57.9, 97.89, 113.79, 120.69, 128.44, 135.64, 136.43, 154.77, 158.01, 160.68; IR (KBr, cm<sup>-1</sup>): 3483, 3255, 3113, 2193, 1643, 1602 (Maleki *et al.*, 2016).

### 3.1.3. Physical data of synthesized pyranopyrazoles

**Table 1.** Physical analytic data of synthesized compounds 5a and 5b

Compound	Structural Formula	Aspect	Color	R <sub>f</sub>	m.p.(°C)	Yield (%)
5a			White	0.36	215-218	87
5b			White	0.65	210-212	98

## 3.2. Toxicological assays

### 3.2.1. Strains and growth conditions

During this study, we screened the effect of the molecule on the protozoan viability to find the toxic lethal

concentration. Accordingly, the results gathered in table 2 demonstrate that both products can be used at their non-toxic concentration (625µg/ml). At this concentration, the growth and viability of this protozoan are not affected.

**Table 2.** The evaluation of Tetrahymena response after treatment with pyranopyrazoles

Toxicological data	Compound	10mg/ml	5mg/ml	2.5mg/ml	1.25mg/ml	0.625mg/ml	0mg/ml
Turbidity	5a	+++	++++	++++	+++++	+++++	+++++
	5b	++	+++++	+++++	+++++	+++++	+++++
Density	5a	0.8x10 <sup>5</sup>	1x10 <sup>5</sup>	1.4x10 <sup>5</sup>	1.6x10 <sup>5</sup>	1.8x10 <sup>5</sup>	2x10 <sup>5</sup>
	5b	1.6x10 <sup>5</sup>	1.8x10 <sup>5</sup>	1.9x10 <sup>5</sup>	2x10 <sup>5</sup>	2.2x10 <sup>5</sup>	


\* (+) means the growth of Tetrahymena in the PPYE medium; - means the absence of Tetrahymena growth in the PPYE medium. The four derivatives are added to the PPYE medium at a non-toxic concentration of 1.25mg/ml. The number of cells is expressed as a function of cells/ml.

### 3.2.2. Treatment with pyranopyrazoles

#### 3.2.2.1. Effect on the growth of Tetrahymena

By comparing and analyzing the results reported in figure 3, we noticed that the supplementation with 5a and 5b derivative protected the normal growth of the protozoan


especially against the oxidative stress; besides that, we found that the compound substituted with a hydroxyl group 5a and 5b are more active against oxidative stress (Figure 3). This effect is more pronounced and appreciated after 48 h of continuous culture.



**Figure 3.** The anti-stress effect of pyranopyrazoles derivatives. Protozoan growth was determined by measuring the absorbance at 600 nm during 120 h. (A) reflected the protective effect of **5a** derivative against oxidative stress and (B) against nitrosative stress.

3.2.2.2. Morphological behavior

Light microscopic analysis of treated cells in the PPYE medium showed clearly that *T. Thermophila* exposed to stress are characterized with morphological transformations such as elongated or spherical forms, the close observation demonstrates the presence of conjugation forms, as well as small dark vacuoles that appeared and increased in number with time. After the treatment of samples with **5a** and **5b** derivatives the protist keeps and preserves their normal shape (pear-shape) against stress damage (see figure 4).



**Figure 4.** Microscopic analysis images of *Tetrahymena thermophila* (TT) cells. Microscopic images were taken at the objective (x10) of cells grown at 32°C in PPYE medium containing H<sub>2</sub>O<sub>2</sub> or SNP as stressors. *Tetrahymena* cells cultivated without stress agents present negative control.

3.2.2.3. The speed swimming of Tetrahymena

The lowest mobility of *Tetrahymena* is recorded under the stress conditions described previously. In close observation, we noticed that some cells moved but not in

stress. The curve (C) represents the protective effect of **5b** derivative against oxidative stress and (D) against nitrosative stress.

straight light as the normal shapes. The protozoans exposed to stress are spinning around themselves. Accordingly, an improvement in speed swimming is observed after the addition of the derivative **5a** and **5b** product, which confirms their protective effect (table 3).


**Table 3.** The protective effect of pyranopyrazoles on physiological parameters of Tetrahymena.

\*The density of *Tetrahymena* cells have been detected by using the light microscope and Malassez chamber.


3.2.2.4. The cells densities

Tetrahymena	Density*	Morphology	Speed swimming(motility)
TT	26.10 <sup>4</sup> cell/ml	Pear-shaped	Rapid moving
TT+SNP	12.10 <sup>4</sup> cell/ml	Atypical form	Slight moving
TT+H2O2	10.10 <sup>4</sup> cell/ml	Weird form	Slight moving
TT+5a+SNP	13.10 <sup>4</sup> cell/ml	Atypical form	Slight moving
TT+5b+SNP	14.10 <sup>4</sup> cell/ml	Atypical form	Slight moving
TT+5a+H <sub>2</sub> O <sub>2</sub>	25.10 <sup>4</sup> cell/ml	Pear-shaped	Rapid moving
TT+5b+H <sub>2</sub> O <sub>2</sub>	20.10 <sup>4</sup> cell/ml	Pear-shaped	Rapid moving

The results reported in Figures 5 and 6 highlighted that the toxic effect of oxidative stress has reduced a significant number of cells, which confirms that *T. Thermophila* is more sensitive to this agent compared to SNP. The supplementation with pyranopyrazoles to the cultures has preserved a large number of cells. The highest activity was recorded after the addition of the **5a** derivative that protected a high number of *Tetrahymena* cells. As reported in Figures 5 and 6, the cell number of treated samples became similar to the control, which proved the protective potential of these compounds.




**Figure 5.** Relative survival of *T. Thermophila* based on cell densities. (A) reflected the protective effect of **5a** derivative against oxidative stress and nitrosative stress(B).




**Figure 6.** Relative survival of *T. Thermophila* based on cell densities. (A) The protective effect of **5b** derivative against oxidative stress and (B) against nitrosative stress. Error bars reflected the standard deviations for three independent biological replicates, with each experiment being comprised of three individual measurements.

### 3.2.3. Antioxidant activity in-vitro: test DPPH

By analyzing and comparing the ongoing results we emphasize that the antioxidant activity of the selected pyranopyrazoles **5a** showed an activity that can exceed the antioxidant activity of the ascorbic acid (see figure 7). By analyzing the data reported in these graphs we noticed that the IC<sub>50</sub> value of **5a** is detected at 6  $\mu\text{g/ml} \pm 0.05$  vs. 9  $\mu\text{g/ml} \pm 0.05$  for ascorbic acid. In parallel, the antioxidant activity of **5b** is important and shows also a significant activity compared to the antioxidant potential of the reference (see figure 8). Indeed, the IC<sub>50</sub> value is recorded at 700  $\mu\text{g/ml} \pm 0.05$  for **5b** compared to 500  $\mu\text{g/ml} \pm 0.05$  for ascorbic acid.



**Figure 7.** Antioxidant activity (% inhibition of DPPH) under different concentrations of **5a** derivative. Determination of IC<sub>50</sub> of the pyranopyrazoles by DPPH method. Ascorbic acid was used as standard.



**Figure 8.** Evaluation of antioxidant properties of pyranopyrazoles **5b** by DPPH assay.

## 4. Discussion

This paper presents a preliminary study describing the protective effect of pyranopyrazoles products against oxidative and nitrosative stress on *Tetrahymena thermophila* growth, to investigate the possible use of these compounds as anti-stress agents. The evaluation of the acute toxicity of both stressors (H<sub>2</sub>O<sub>2</sub> and SNP) on *Tetrahymena* cells will be considered as an important reference value for its further potential utilization to control chronic diseases. This protozoa *Tetrahymena* is characterized by three essential phases of growth (Errafiy and Soukri, 2012): a latent phase that lasts 24 h followed by an exponential phase and a stationary phase that approaches 160h (Mar, El khalfi and Soukri, 2018). When protozoa are treated with stressors, there is a disruption in protozoan growth that varies with the species and stress agent used (Mountassif *et al.*, 2007).

Our finding confirmed that Hydrogen peroxide completely inhibits the growth of *Tetrahymena Thermophila* however the nitroprusside sodium slows its growth (Mar, El khalfi and Soukri, 2018). Similar results have been reported previously by Errafiy (Errafiy, Ammar, and Soukri, 2013). Some biological and molecular mechanisms in responses to stress have been examined by using many model organisms including protozoan, yeasts, and vertebrates (Errafiy and Soukri, 2012).

The Toxicities of stressors are routinely determined by analysis of some parameters such as the dynamic growth curves of *T. Thermophila*; the ciliary movement and the viability of the cells (Addoum *et al.*, 2018). In this study,

the authors reported that the treated protozoan with peroxide hydrogen undergo some morphological and physiological transformations, and according to the literature surveys the Hydrogen peroxide acts in oxidative stress as a messenger molecule that diffuses through cells to initiate intermediate cellular effects (Ganjoor and Mehrabi, 2017) such as changes in the shape and recruitment of immune cells (Rahman *et al.*, 2012). Prompted by our finding, we confirmed that the cell densities of the *Tetrahymena* population were proportional to the IC50 concentration of stressors. At this lethal concentration, cell growth diminished to 50 %. Similar results were observed also by using other stressors such as ethanol (Nilsson, 1974), bromide ethidium (Ashour B, 1980), gelsemine (Ye *et al.*, 2019), and some xenobiotics. These stressors may influence a variety of physiological parameters such as the membrane fluidity, autophagic vacuolization, the rates of cell death and the main speed of *Tetrahymena* cells (Fryb and Henry, 1997)

Nitrosative stress, as well as oxidative stress, can activate a specific route of signalization via the overproduction of NO; this radical can react with ROS such as superoxide anion ( $O_2^{\cdot-}$ ) to form peroxyntirite ( $ONOO^-$ ) (Sies, 2017). The peroxyntirite is recognized as a biological oxidant that influenced some mitochondrial functions and induced cell death by apoptosis (Sies, 2017).

In our study, we induced nitrosative stress by using the SNP, as a consequence we detected several morphological changes including the appearance of mostly elongated or spherical forms with wide gloomy vacuoles. All that, reflected the harmful effect of SNP that can damage the structure and alter some cellular constituents of the stressed protozoan (DNA; mitochondria; membrane...). On the other hand, excessive NO production can induce a reversible (or irreversible) modification of cellular proteins (Fourrat *et al.*, 2007). NO-induced stress inhibits several enzymes essential for metabolism, including the specific activity of GAPDH via S-nitrosylation correlated with a significant decrease in the growth of protozoan *Tetrahymena Thermophila*. By comparing the effect of both derivatives on the morphology and growth of protozoa in the presence of stressors, we can deduce that **5a** and **5b** compounds have a higher protective effect against the  $H_2O_2$ -induced stress. The main speed of *Tetrahymena* (motility) is another physiological parameter which should be more studied to assess the cytotoxic effect of stress on the locomotion of this ciliate. The ciliary locomotion of treated protozoan is diminished, which can be an indicator of membrane toxicity; it can be also explained by an alteration of membrane potency. Several research studies indicate that toxicants alter the ion fluxes which have a direct effect on the swimming speed of *Tetrahymena* (Fryb and Henry, 1997)

To the best of our knowledge, the current study is devoted to examining the protective effects of pyranopyrazoles scaffold against stress-induced *Tetrahymena* cells. The supplementation with **5a** and **5b** compound has been improved the motility, the cell number, and the growth curve; which emphasizes the protective effect of both derivatives especially against oxidative damage (Mar *et al.*, 2019). We also notified that the **5a** derivative is the most active compound that stimulates also the growth of *Tetrahymena*; this effect can be related to their basic pyranopyrazole skeleton of this

compound and the nature of the substituent on the 4-phenyl ring. To remedy the damage caused by oxidative and nitrosative stress, we used two pyranopyrazoles derivatives. These compounds are characterized by several properties, such as antimicrobial, antitumoral, and antioxidant (Shahbazi *et al.* 2019). Nevertheless, these synthesized compounds own a toxic effect against the protozoa *Tetrahymena*. The non-lethal concentration of the two derivatives is used to evaluate their protective effect against stressors. The results reported herein showed clearly that **5a** and **5b** protect protozoa against extracellular  $H_2O_2$ , which can be related to the increase of SOD and glutathione peroxidase production (Mountassif *et al.*, 2007). According to the literature surveys, these synthesized products were reported to have significant antioxidant activity. For this purpose, we can say that both compounds used in our study can have a direct effect on the activity of ROS by trapping them or an indirect effect by increasing the production of intracellular antioxidant enzymes (Mar *et al.*, 2019).

On the other hand, the study in vitro results reveal that **5a** derivative possesses the top of antioxidant activity compared to the antioxidant of reference. The IC 50 of this product is lower than the value of ascorbic acid ( $6 \mu\text{g mL}^{-1}$  for **5a** vs  $9 \mu\text{g mL}^{-1}$  for Ascorbic acid). The second product **5b** besides also a good activity (IC<sub>50</sub> value is recorded at 7mg/ml) compared to ascorbic (IC<sub>50</sub> value is 5mg/ml). Overall, our finding established an interesting structure-activity relationship SAR. Likewise, the compound incorporating the hydroxyl group in the *meta*-position of the phenyl group (derivative **5a**) provided an impressive activity against oxidative stress. The most illustrative example has been reported by (Roghayeh, and Mahmoodi 2016), they indicated that the introduction of a hydroxyl group on the aromatic ring of pyranopyrazole can effectively modulate its mechanism of action and increase significantly their free radical scavenging activities (Jouha *et al.*, 2017; Harshad G *et al.*, 2012). These results are in harmony with our results reported herein.

In summary, the pyranopyrazoles are highly active compounds. Although its chemical structure and stereochemistry have been defined (Reddy *et al.*, 2018), nevertheless its biological effects are poorly described.

## 5. Conclusion

We described herein a simple and efficient approach to synthesize some pyranopyrazoles derivatives. The results reported in this paper confirmed their antioxidant potential. We are convinced that the current study is a step to underlying the relationship between structure and biological function which will encourage to building-up a variety of motifs occurred via structural modifications onto existing bioactive pyranopyrazoles scaffolds, as a consequence we may enhance the inherent medicinal deed of these compounds or provide their drug-like proprieties.

## Acknowledgment

Financial support by the National Center for Scientific and Technical Research (CNRST) is gratefully acknowledged. This work was supported also by University Hassan II of Casablanca. All authors wish to

thank Professor Aziz AUHMANI for NMR analysis in the Faculty of Sciences Cadi Ayyad Marrakech in Morocco.

### Conflict of interest

All the authors declared that there is no conflict of interest.

### Funding statement

This research did not receive any specific grant from funding agencies in the public, commercial, or non-profit sectors.

### References

- Addoum B, El Khalfi B, Mar PD, Filali Ansari Na, Elmakssoudi A, Soukri A. 2018. Synthesis, Characterization of Some  $\alpha$ -Aminophosphonate Derivatives and Comparative Assessment of Their Antioxidant Potential in-Vivo and in-Vitro. *Der Pharma Chemica*, **10(12)**:58-67.
- Ashour B, Tribe M, and Whittaker P. 1980. The effect of chloramphenicol, ethidium bromide and cycloheximide on mortality and mitochondrial protein synthesis of adult blowflies. *Journal of Cell Science*, **41**:273-289.
- Azaam M, El Refaie K, Bad A, El-din A, Khamis A, and El-Magd M. 2018. Antioxidant and Anticancer Activities of  $\alpha$ -Aminophosphonates Containing Thiadiazole Moiety. *Journal of Saudi Chemical Society*, **22(1)**:34-41.
- Debabis D, Banerjee R, and Mitra A. 2014. Bioactive and Pharmacologically Important Pyrano[2,3-c]Pyrazoles. *Journal of Chemical and Pharmaceutical Research*, **6(11)**:108-116.
- Doerder FP and Brunk C. 2012. Learn More about *Tetrahymena Thermophila*. The *Zebrafish: Genetics, Genomics and Informatics*:30-50.
- Dove P. 2016. Docking Studies of Hybrid Pyrazole Analogues. *Drug Design, Development and Therapy*, **10**: 3529-3543.
- El-borai M, Rizk HF, Sadek M, and El-keiy M. 2016. An Eco-Friendly Synthesis of Heterocyclic Moieties Condensed with Pyrazole System under Green Conditions and Their Biological Activity. *Green and Sustainable Chemistry*, **06(02)**:88-100.
- Errafiy N, Ammar E and Soukri A. 2013. Protective Effect of Some Essential Oils against Oxidative and Nitrosative Stress on *Tetrahymena Thermophila* Growth. *Journal of Essential Oil Research*, **25(4)**:339-47.
- Errafiy N and Soukri A. 2012. Purification and Partial Characterization of Glyceraldehyde-3-Phosphate Dehydrogenase from the Ciliate *Tetrahymena Thermophila*. *Acta Biochimica Et Biophysica Sinica*, **44(6)**:527-34.
- Fourrat L, Iddar A, Valverde F, Serrano A, and Soukri A. 2007. Effects of Oxidative and Nitrosative Stress on *Tetrahymena Pyriformis* Glyceraldehyde-3-Phosphate Dehydrogenase. *The Journal of Eukaryotic Microbiology*, **54(4)**:338-46.
- Fryb H and Henry JA. 1997. Membrane Toxicity of Opioids Measured by Protozoan. *Toxicology*, **117(1)**:35-44.
- Ganjoor M and Mehrabi M. 2017. Inhibitory Effect of Hydrogen Peroxide and Ionic Silver on Growth of a Pathogenic Bacterium (*Vibrio Harveyi*) Isolated from Shrimp (*Litopenaeus Vannamei*). *Fish & Ocean Opj*: 2(4).
- Harshad G, Ranjan G and Manish P. 2012. Microwave-Assisted Multi-Component Synthesis of Indol-3-Yl Substituted Pyrano[2,3-c] Pyrazoles and Their Antimicrobial Activity. *Journal of the Serbian Chemical Society*, **77(8)**:983-91.
- Huang AG, Xiao T, Lei L, Gao X W, and Fei L. 2016. The Oxidative Stress Response of Myclobutanil and Cyproconazole on *Tetrahymena Thermophila*. *Environmental Toxicology and Pharmacology*, **41**:211-18.
- Jouha J, Loubidi M, Bouali J, Hamri S, Hafid A, Suzenet F, Guillaumet G, Dagci T, Khouili M, Aydin F, Saso L, and Armagan G. 2017. Synthesis of New Heterocyclic Compounds Based on Pyrazolopyridine Scaffold and Evaluation of Their Neuroprotective Potential in MPP+-Induced Neurodegeneration. *European Journal of Medicinal Chemistry*, **129**:41-52.
- Kanagaraj K and Pitchumani K. 2010. Solvent-Free Multicomponent Synthesis of Pyranopyrazoles: Per-6-amino-bcyclodextrin as a Remarkable Catalyst and Host. *Tetrahedron Letters*, **51** :3312-3316.
- Kumar KA, Verma AK, Kumar T, and Srivastava R. 2015. Oxidative Stress: A Review. *The international journal of science & Technologie*, **3(7)**: 2321 - 919.
- Mamaghani M, and Hossein Nia R. 2019. A Review on the Recent Multicomponent Synthesis of Pyranopyrazoles. *Polycyclic Aromatic Compounds*.
- Maleki B, Nasiri N, Reza T, Khojastehnezhad A, and Akhlaghi HA. 2016. Green Synthesis of Tetrahydrobenzo[b]Pyrans, Pyrano[2,3-c] Pyrazoles and Spiro[Indoline-3,4'-Pyrano[2,3-c]Pyrazoles Catalyzed by Nano-Structured Diphosphate in Water. *RSC Advances*, **6** :79128-79134.
- Mar PD, Otto H, El khalif B, and Soukri A. 2019. Biochemical Study of the Protective Effect of *Salvia Officinalis* Essential Oil against the Oxidative Stress Induced by Hydrogen Peroxide in *Tetrahymena Thermophila*. *Derpharma chemica*, **11(3)**:8-12.
- Mar PD, El Khalfi B, and Soukri A. 2018. Protective Effect of *Oregano* and *Sage* Essentials Oils against Oxidative Stress in *Tetrahymena Thermophila* and *Tetrahymena Pyriformis*. *Journal of King Saud University - Science*, **32** :279-287.
- Mar PD, El Khalfi B, Perez-Castiñeira JR, Serrano A, Soukri A. 2017. Cell Stress by Phosphate of Two Protozoa *Tetrahymena Thermophila* and *Tetrahymena Pyriformis*. *Advances in Bioscience and Biotechnology*, **8**:451-62.
- Mountassif D, Kabine M, Manar R, Bourhim N, Zaroual Z, Latruffe N, and El Kebbj MS. 2007. Physiological, Morphological and Metabolic Changes in *Tetrahymena Pyriformis* for the in Vivo Cytotoxicity Assessment of Metallic Pollution: Impact on d- $\beta$ -Hydroxybutyrate Dehydrogenase. *Ecological Indicators* **7(4)**:882-94.
- Nilsson JR. 1974. Effects of DMSO on Vacuole Formation, Contractile Vacuole Function, and Nuclear Division in *Tetrahymena Pyriformis* GL. *Journal of Cell Science*, **16(1)**:39-47.
- Pisoschi AM, and Pop A. 2015. The Role of Antioxidants in the Chemistry of Oxidative Stress: A Review. *European Journal of Medicinal Chemistry*, **97**:55-74.
- Rahman T, Hosen I, Towhidul IM, and Shekhar HU. 2012. Oxidative Stress and Human Health. *Advances in Bioscience and Biotechnology*, **03(07)**:997-1019.
- Reddy GM, Jarem RG, Reddy VH, KumariAK, ZyryanovG V, and YuvarajaG. 2019. An Efficient and Green Approach: One Pot, Multi Component, Reusable Catalyzed Synthesis of Pyranopyrazoles and Investigation of Biological Assays. *Journal of Saudi Chemical Society*, **23(3)**:263-73.
- Reddy GM, Jarem RG, Reddy VH, KumariAK, ZyryanovG V, and YuvarajaG. 2018. Highly Functionalized Pyranopyrazoles: Synthesis, Antimicrobial Activity, Simulation Studies and Their Structure Activity Relationships (SARs). *Research on Chemical Intermediates*, **44(12)**:7491-7507.



- Roghayeh SM. 2016. Green and efficient Synthesis of Pyranopyrazoles Using [ Bmim] [ OH À ] as an Ionic Liquid Catalyst in Water under Microwave Irradiation and Investigation of Their Antioxidant Activity. *RSC Advances*: 85877–84.
- Shahbazi S, Ghasemzadeh MA, Pegah S, Zolfaghari MR, and Bahmani Ma. 2019. Synthesis and Antimicrobial Study of 1,4-Dihydropyrano[2,3-c] Pyrazole Derivatives in the Presence of Amino-Functionalized Silica-Coated Cobalt Oxide Nanostructures as Catalyst. *Polyhedron*, **170**:172–79.
- Sies H. 2017. Redox Biology Hydrogen Peroxide as a Central Redox Signaling Molecule in Physiological Oxidative Stress: Oxidative Eustress. *Redox Biology*, **11**:613–19.
- Sugimoto N, Watanabe H, and Ide A. 1960. The Synthesis of L- $\alpha$ -Amino- $\beta$ -(Pyrazolyl-N)-Propionic Acid in *Citrullus Vulgaris*. *Tetrahedron*, **11**(4):231–33.
- Tekale SU, Kauthale SS, JadhavKM, and Rajendra P P. 2013. Nano-ZnO Catalyzed Green and Efficient One-Pot Four-Component Synthesis of Pyranopyrazoles. *Journal of Chemistry*, **840954** :8.
- Ye Q, Zhang C, Wang Z, Feng Y, Zhou A, Shaolin X, Qiong X, Enfeng S, and Jixing Z. 2019. Induction of Oxidative Stress, Apoptosis and DNA Damage by Koumine in *Tetrahymena Thermophila*. *PLoS ONE*, **14**(2): e0212231
- Zahouily M, Elmakssoudi A, Mezdar A, Rayadh A, Sebti S, and Lazrek H. 2005. Three Components Coupling Catalysed by Na<sub>2</sub>CaP<sub>2</sub>O<sub>7</sub>: Synthesis of Amino Phosphonates Under Solvent-Free Conditions at Room Temperature. *Letters in Organic Chemistry*, **2**(5):428–32.
- Zhang SW, Feng JN, CaoY, Meng LP, and Wang SL. 2015. Autophagy Prevents Autophagic Cell Death in *Tetrahymena* in Response to Oxidative Stress. *Zoological Research*, **36**(1): 167-173.