

Embryonic development of the striped spiny eel, *Mastacembelus pancalus* (Hamilton, 1822) in captive condition

Md. Sherazul Islam* and Tandra Rani

Department of Fisheries and Marine Bioscience, Jashore University of Science and Technology, Jessore-7408, Bangladesh

Received: May 8, 2020; Revised: October 8, 2020; Accepted: Oct 26, 2020

Abstract

The striped spiny eel, *Mastacembelus pancalus* is one of the common food fish in Bangladesh, but lacking study on embryonic development of the fish. The present study describes the embryonic progress stages of the fish in the confined state. The embryogenesis is divided into seven major phases: Zygote phase, Cleavage phase, Morula phase, Blastula phase, Gastrula phase, Organogenesis phase and Hatching phase. According to distinct development characteristics, the further sub-stages were also developed whenever possible. Fertilized eggs were sticky, colorless, and demersal with identical perivitelline space. The thickness of the unfertilized eggs ranged from 0.55-0.58 mm, which increased to 0.62-0.71 mm in fertilized eggs. The perivitelline space development happened at 0.35 h after fertilization (AF). The first cleavage groove occurred at the animal pole (two cells) 0.55 h AF. The added cell division such as four cells, eight cells, sixteen cells, thirty-two cells, and multi cells stages was initiated at 1.25, 1.45, 2.15, 2.50, and 4.40 h AF respectively. The morula, blastula, gastrula, yolk plug stage, and organogenesis stages were seen at 8.25, 11.30, 17.15-23.50, 29.20, and 34.15-35.30 h AF respectively. The initial heartbeat was observed at 34.00 h AF. The head and tail ends of the embryo were distinguished at 35.00 h AF. The notochord was also evident at the same hours. The embryo started to hatch at 36.00 h AF which accomplished 39.00 h at 29.11 ± 0.29 °C. The new hatchling was 1.65 ± 0.15 mm in average length. The present findings will serve as baseline information to develop the breeding protocol of the species in the hatchery condition.

Keywords: Embryogenesis, cleavage, organogenesis, hatching

1. Introduction

The Guchibaim (*Mastacembelus pancalus*) is one of the regular food fish found in Asian countries, namely in India, Pakistan, Bangladesh, and Nepal (Talwar and Jhingran, 1991; Froese and Pauly, 2006) and recognized as a striped spiny eel. The species is a common and demandable food fish in Bangladesh and is locally known as 'guchi baim.'

In the past, the fish was available in estuaries and freshwater habitats throughout Bangladesh (Ali, 1967). With the destruction of natural habitat including overexploitation, the fish has diminished abruptly from wildlife (Afroz *et al.*, 2014). Besides drying up of downcast land and using pesticides, the ordinary production lands of this fish are in threat (Rahman *et al.*, 2009). Besides these, the fish is collected only from nature that exaggerates the natural reduction of the fish.

The fish is critically endangered in Bangladesh (FISHWISE, 2013) but has not been shown in the red list of IUCN (Anonymous, 2006). It is needed for the management of natural habitat or to introduce artificial propagation as well as culture. Thus, it is essential to realize the embryonic development stages of the fish to set up a non-natural propagation in a confined condition, it's stocking at mass scale for its expansion and conservation.

Study on biology and breeding has been done on different eel fishes such as on *Mastacembelus pancalus* (Hasan *et al.*, 2016; Karim and Hossain, 1972); *M. armatus* (Serajuddin and Mustafa, 1994); *Macrogathus aculeatus* (Das and Kalita, 2003); *M. pancalus* (Suresh *et al.*, 2006). The development of egg and/or larvae on eel fishes like *M. pancalus* (Rahman *et al.*, 2009; Afroz *et al.*, 2014), *M. aculeatus* (Sahoo *et al.*, 2007; Farid *et al.*, 2008); *Muraenesox cinereus* (Umezawa *et al.*, 1991); *Anguilla rostrata* (Oliveira and Hable, 2010); *Mastacembelus mastacembelus* (Sahinoz *et al.*, 2006) have done, but there are no details of basic work on embryogenesis of *M. pancalus* except Rahman *et al.* (2009). As the embryonic development process differs from species to species, it is important to know the detailed developmental stages of any fish to consider the artificial propagation of the fish in captivity. It is an urgent need to develop a captive breeding protocol of the species due to the declination of natural propagation. It is essential to know the variations of features in embryonic development and to know the development of organs for the management and rearing technology for seed production of any fish species. Thus, detailed embryonic developmental stages with prominent features were carried out in the captive condition of *M. pancalus*.

* Corresponding author e-mail: dms.islam@just.edu.bd.

** **Abbreviations** : AFAfter fertilization; EM Egg membrane ; VM Vitelline membrane ; PS Perivitelline space

2. Materials and Methods

2.1. Study site and preparation of aquarium

The experiment was carried out in the laboratory of the Department of Fisheries and Marine Bioscience, Jashore University of Science and Technology, Jashore, Bangladesh. The experiment was done in rectangular glass aquaria (3 f length \times 1.16 f width \times 1.25 f depth), each containing thirty liters of water. Water hyacinths were used as a source of plants, and their roots were used as a substrate to lay the eggs of the fishes. The major physicochemical parameters like temperature, pH, and DO were measured at each two hours interval during embryonic development. A pH meter (EZODO, 7200, Taiwan) and a DO meter (LTLutron YK-22DO, Taiwan) were used to measure the water pH, DO and temperature respectively.

2.2. Collection of egg

In the same laboratory, eggs of *M. pancalus* were produced by inducing with PG hormone. Brood fishes were collected from the nearby natural habitat (Baor) and the average body weight was 9 g of each. There were three treatments, and each had two more replications for the inducing of the fish. Three pairs of broods were kept 1:1 (male: female) the ratio in each aquarium for natural propagation after inducing. Fishes were spawned (100%) within sixteen to twenty hours after the administration of the hormone. The sticky eggs of the *M. pancalus* were attached to the aquatic weeds and sometimes on the substrates. Eggs were sampled softly either along with the roots of the water hyacinth or using a father in each sampling.

2.3. Observation of embryonic development

The embryonic development stages were observed, and a snap was taken with a photographable microscope (Carl Zeiss microscopy GmbH, S.N. MKG8639, Germany). The eggs were observed at every 5 to 10 minutes interval till the accomplishment of the morula and then after observed each one-hour intermission until hatching. The diameter of oocytes and eggs was measured at each sampling time using the microscopic camera. The development stages and characteristics were confirmed by following Sahoo *et al.* (2007), Rahman *et al.* (2009), and Honji *et al.* (2012).

3. Results

3.1. Aquarium environment

Water parameters play a major role in the spawning, embryonic development, and hatching of any fish species. In the present study, the physicochemical condition of spawning aquaria such as temperature, dissolved oxygen, and pH ranged from 28.8 to 29.4 °C, 5.25 to 5.75 mg.L⁻¹ and 7.75 to 8.12 respectively.

3.2. Characteristics of the egg

The yellowish eggs were spherical, demersal, and adhesive. They stuck to the roots of water hyacinth in the aquarium. The unfertilized eggs were opaque while the fertilized eggs were transparent with visible egg membrane and yolk (Figure 1A, 1B). The diameter of the fertilized eggs increased 0.62 to 0.71 mm from 0.55 to 0.58 mm of the unfertilized egg.




Figure 1. The unfertilized (A) and fertilized (B) egg of *M. pancalus*. The formation of perivitelline space (C) at the early stage of the zygote. EM = Egg membrane; VM = Vitelline membrane; PS = perivitelline space; OG = oil globule.

3.3. Embryonic development

After the fertilization of eggs, the embryonic period started and ended at the time of accomplishment of the general organ systems is common in all fishes. The

fertilized eggs hatched out within 36 to 39 h after fertilization (AF). The events in embryonic progress and their respective time and features of *M. pancalus* are presented in table 1 and figure 1-4.

Table 1. Embryonic developmental stages (major and sub-phases) with the respected time and features of *M. pancalus* in aquarium condition.

Major phase	Sub-phase	Figure	Time (h/min)	Developmental features
Zygote	Fertilized eggs	1	0.00	Eggs adhesive, demersal, spherical, and transparent.
	Perivitelline space formation		0.25- 0.30	Formation of perivitelline space around the yolk
Cleavage	Blastodisc	2	0.35	Formation of blastodisc at the animal pole
	Two cell		0.40-0.50	Aggregated oil globules at the animal pole, the commencement of the first cleavage
	Four cell		1.25	2 nd cleavage, 4-cell
	Eight cell		1.25-1.45	The blastomeres were unequal in size and remain in two rows. 8-cell
	Sixteen cell		2.15	4 th cleavage, 16-cell
	Multi-cell		3.00 -4.40	Quick successive division and transformed into 32, 64, 128 celled stage and so on
Morula			7.15-8.25	Formed a cap at the animal pole, which gradually increased in size.
Blastula		3	11.30-14.45	The marginal blastomeres lost their boundaries and were compressed.
Gastrula	Early gastrula		17.15	Blastoderm started to form a thin layer by invading the yolk and overthrowing over the yolk
	Middle gastrula		22.10	Development of germinal ring in the region of yolk and about ½ of the yolk was possessed by blastoderm.
	Late gastrula		23.50	The embryonic shield was visible and the blastoderm covered ¾ of the yolk.
Organogenesis	Embryonic body		27.15	The embryonic body was visible.
	Yolk plug		29.20	Finished yolk invasion, visible undeveloped head and tail and became differentiated.
	Segmentation	4	34.15- 35.35	Distinguished head and tail, and notify of beating heart. Noticeable notochord in cellular structure.
Hatching			36.00-39.00	The twisting movement became more forceful and the embryo broke the egg pod.

3.3.1. Zygote phase

This phase is characterized by fertilized and perivitelline space formation. The perivitelline space formation occurred at 0.30 h AF in the present study. The perivitelline space (the thin space that separated the egg membrane) was fluid-filled and equal all around the egg membrane (Figure 1C). Oil globules were visible in the yolk in this phase.

3.3.2. Cleavage phase

The single-cell stage became clear with the accumulation of cytoplasm over the animal pole as a protrusion at 0.35 h AF representing the early blastodisc or germinal disc stage (Figure 2A). As the development proceeded, the oil globules were found to aggregate and the cytoplasmic disc became thick and the first cleavage

occurred within 1.00 to 1.25 h AF. The vertical cleavage occurred which divided the blastodisc into two different cells at 0.50 h AF (Fig. 2B). The further cleavage was at a right angle to the first and observed forming four cells within 1.25 h (Figure 2C). Further division of blastomere took place with the advancement of time to reach eight-cell and sixteen cell stages at 1.45 h and 2.15 h AF respectively (Figure 2D, 2E). After quick succession, the sixteen-celled stages resolved into 32, 64, 128 celled stages, and so on. However, due to the rapid occurrence of these cell divisions, it was not possible to observe or count the stages; and hence in the present study, it was considered as a multi-celled stage (Figure 2F). Eggs were measured and noticed the same size (0.62 – 0.71 mm).




Figure 2. The cleavage phase of *M. pancalus*; Formation of blastodisc (A), 2-cell (B), 4-cell (C), 8-cell (D), 16-cell (E), and Multi-cell stage (F). GD=Germinal disc (blastodisc); OG=Oil globule.

3.3.3. Morula phase

The blastomeres were reduced in size and accumulated around the animal pole during the morula stage. A cap-like creation was seen at the animal pole, which size increased slowly (Figure 3A). The morula phase was recognized at 8.25 h AF.

3.3.4. Blastula phase

The embryo was further divided into numerous cells after the morula and formed a blastoderm by arranging a form of a layer (Figure 3B). The blastodisc was formed by the gradual formation of several layers due to further cell division. At this phase, blastocoels also appeared (a space between yolk and blastoderm). This phase of the embryo is

called 'blastula' which was observed within 11.30 h to 14.45 h AF.

3.3.5. Gastrula phase

The gastrulation phase is subdivided into three stages: like early gastrula, middle gastrula, and late gastrula. The incursion of the yolk started by blastoderm through spreading over the yolk like a thin layer which is denoted as the early gastrula and resulted within 17.15 h AF (Figure 3C). In the middle of gastrulation, noticed a visible germinal ring on every side of the yolk. In this stage, about half of the yolk was engaged by blastoderm (Figure 3D). The embryonic shield was visible at late gastrulation and blastoderm covered $\frac{3}{4}$ th of the yolk (Figure 3E).




Figure 3. The morula (A), blastula (B) and gastrula phase (C, D, E) of *M. pancalus*; C=Early gastrula, D=Middle gastrula and E=Late gastrula.

3.3.6. Organogenesis phase

The embryonic body formation appeared at 27.15 h AF (Figure 4A). The gradual spreading above the germ layer in the plug stage completed the yolk incursion. The head and tail seemed in this stage within 29.20 h AF (Figure

4B). In addition to this, the embryo was lengthened and bordered the yolk materials and differentiated the tail and head ends (Figure 4C). The first beating heart was visible at about 34.00 h AF. Within the cellular structure, the notochord became noticeable within 34.15 to 35.35 h AF.




Figure 4. The organogenesis and hatching of *M. pancalus*; Formation of the embryonic body (A), Yolk plug stage (B), segmentation of organ (C), and just before hatching (D).

3.3.7. Hatching phase

In this phase, noticed elongated embryo which progressively separated. The tail became steadily separated from the yolk mass (Fig. 4D). The embryo started an irregular twisting movement. Later, the eggshell started to rupture by the embryos due to the continuous movement. The hatch out of larvae was noticed in 36.00 to 39.00 h after fertilization with its tail portion first and completed within 3.10 h. The size of the newly hatched larvae was 1.65 ± 0.15 mm in length (Fig. 5).




Figure 5. The new hatchling of *M. pancalus* in aquarium condition. Hatching started at 36 h after fertilization.

4. Discussion

4.1. Aquarium environment

The embryonic development of fishes is directly related to the water parameters, particularly water temperature. In the present study, the recorded water parameters were deemed suitable for the species. Water temperature during the time of embryonic development was within 28.8 to 29.4 °C which was close to other studies like 28 to 29 °C (Sahoo *et al.*, 2007) and 27-31 °C (Rahman *et al.*, 2009).

4.2. Characteristics of the egg

The characteristics of the fertilized and unfertilized eggs are more or less similar to other studies. The unfertilized eggs were dense, demersal, sticky and the fertilized eggs were round, clear, and sticky which is supported by Rahman *et al.* (2009). Though the same characteristics of fertilized eggs shown in

Mastacembelidae, different egg color such as 'green color' was observed in the case of *M. aculeatus* (Sahoo *et al.*, 2007). This variation may be due to species variation. The fertilized egg diameter in the present study recorded was 0.62 to 0.71 mm which was smaller than what was reported by Rahman *et al.* (2009). They recorded up to 0.70 to 1.30 mm the size of the fertilized egg in *M. pancalus*. However, further larger (1.20 to 1.40 mm) reported by Sahoo *et al.* (2007) in *M. aculeatus* and 1.50 to 2.02 mm in *M. mastacemblus* reported by Sahinoz *et al.* (2006).

4.3. Embryonic development

The cell division pattern and the further embryonic development stages were more or less similar to other studies of Mastacembelidae. However, differences were noticed in the case of the time of development in different species and even within the same species. The presence of perivitelline space in the present study was also reported in fertilized eggs of New Zealand freshwater eel, *A. dieffenbachia* (Lokman and Young, 2000), and Mesopotamian spiny eel, *M. mastacemblus* (Sahinoz *et al.*, 2006).

The initiation of 1st cleavage, the formation of blastodisc at the animal pole was noticed at 0.35 h AF which was similar to *M. pancalus* (Rahman *et al.*, 2009) and *M. aculeatus* (Sahoo *et al.*, 2007). However, in the case of *M. mastacemblus* it took about 4.00 h (Sahinoz *et al.*, 2006). In the present study, it was noticed that cell division was completed within about 5.00 h AF which was almost the same reported by Rahman *et al.* (2009). However, in the case of *M. aculeatus*, the cell division was completed shortly, and it was by 3.30 h AF (Sahoo *et al.*, 2007). Moreover, within the same species, the induction of morula showed at different times. In the present study, the morula stage appeared within 8.25 h AF whereas Rahman *et al.* (2009) observed this stage at 10.10 h AF. According to Sahoo *et al.* (2007), the same stage in *M. aculeatus* occurred at 4.10 h AF. The blastoderm enclosed nearly 3/4th of the yolk and an embryonic body was formed 27.15 h AF which was noticed 24.30 h AF in the same species (Rahman *et al.*, 2009) and 25.30h AF in *M. aculeatus* (Sahoo *et al.*, 2007) whereas more time (40h) was taken in the case of *M. mastacemblus* (Sahinoz *et al.*, 2006).

The formation of the head and tail of the embryo showed 29.20h AF which was observed in 31.30 h in the same species (Rahman *et al.*, 2009) and even in higher temperatures. However, at a similar temperature, this characteristic was observed at 4.00 h earlier compared to *M. aculeatus* (Sahoo *et al.* (2007) and took more than double duration (77h) in *M. mastacemblus* (Sahinoz *et al.* 2006). The heart pulsation was noticed during about 34.00 h AF in this study alike to Rahman *et al.* (2009) but earlier as compared to Japanese eel, *A. paponica* (Yamamoto *et al.*, 1995). The twisting movement and first hatching were observed at 36.00 h AF. The earlier twisting movement and first hatching were reported by Rahman *et al.* (2009) in the same species and in similar water temperatures. In the case of other eel fishes, different hatching time was reported like 31.45 h in *M. aculeatus* (Sahoo *et al.*, 2007), 38.00-45.00 h in *A. paponica* (Yamamoto *et al.*, 1995) and 85.00 h in *M. mastacemblus* (Sahinoz *et al.*, 2006).

In the present study, it was strong evidence that quick development happens until morula compared to other

studies, but later stages took a longer period particularly in organogenesis and hatching (Figure 6). This may be due to temperature variation as compared to Rahman *et al.* (2009) who recorded more than 30°C after gastrulation whereas in

the present study it was less than 30°C. The other variability like the formation of morula and head and tail due to the different rates of development in different species in addition to temperature.




Figure 6. Major events during the embryonic development of *M. pancalus*. C=completion, F=formation, E=Embryonic

5. Conclusion

The study generated detailed information on early developmental commencement with distinguishing characteristics of *M. pancalus*. In conclusion, it is said that the embryonic development of *M. pancalus* commonly imitates that of other eel fishes. However, the period of development varies in some stages. Besides, the development rate of the embryo varied in the variation of water temperature. The development process is noticed faster in the higher the temperature, and vice versa.

Acknowledgement

The present study was financially supported by the Grant for Advanced Research in Education (GARE), Ministry of Education, People's Republic of Bangladesh. We also thank other groups of the project who successfully breed the species in the aquarium by inducing and providing us with the eggs for further study.

References

- Afroz A, Islam MS, Hasan MR, Hasnahena M and Tuly DM. 2014. Larval rearing of spiny eel, *Mastacembelus pancalus* in the captivity with emphasis on their development stages. *Int J Fish Aquat Stud*, **1(6)**: 163-167.
- Ali MH. 1967. Induced breeding of major carps in ponds by pituitary hormone injection. Agricultural Information Service, Dhaka. pp. 23-26.
- Anonymous. IUCN red list of threatened species (2006). International Union for Conservation of Nature. <http://www.redlist.organization> (January 2019).
- Das SK and Kalita N. 2003. Captive breeding of peacock eel, *Macrognathus aculeatus*. *Aquac Asia*, **8**: 17-21.
- Farid SM, Miah MI, Akter MD, Saha D and Rahman MM. 2008. Embryonic and larval development of tarabaim (*Macrognathus aculeatus*). *J Agrofor Env*, **2(2)**: 123-129.
- FISHWISE, <http://www.fishwise.co.za>, Species=*Mastacembelus pancalus* 5 January, 2013.
- Froese R and Pauly D. (eds), 2006. Fishbase, World-wide-Web Electronic publication. <http://www.fishbase.org>

Hasan MR, Islam MS, Afroze A, Bahdur P and Akter S. 2016. Captive breeding of Striped Spiny Eel, *Mastacembelus pancalus* (Hamilton, 1822) considering the various hormonal responses. *Int J Fish Aquat Stud*, **4(3)**: 07-11.

Honji MR, Tolussi CE, Mello PH, Caneppele D and Moreira RG. 2012. Embryonic development and larval stages of *Steindachneridion parahybae*-implications for the conservation and rearing of this endangered Neotropical species. *Neotrop Ichthyol*, **10(2)**: 313-327.

Karim MA and Hossain A. 1972. Studies on the biology of *Mastacembelus pancalus* (Spiny Eel, Hamilton) in artificial pond. Part II. Sexual maturity and Fecundity. *Bangladesh J Agric Bio Sci*, **1(2)**:15-18.

Lokman PM and Young G. 2000. Induced spawning and early ontogeny of New Zealand freshwater eels (*Anguilla dieffenbachii* and *A. australis*). *NZ J Mar Freshwater Res*, **34**: 135-145.

Oliveira K and Hable WE. 2010. Artificial maturation, fertilization, and early development of American eel (*Anguilla rostrata*). *Can J Zool*, **88**:1121-1128.

Rahman MM, Miah MI, Taher MA and Hasan MM. 2009. Embryonic and larval development of guchibaim, *Mastacembelus pancalus* (Hamilton). *J Bangladesh Agril Univ*, **7 (1)**: 193-204.

Sahinoz E, Dogu Z and Aral F. 2006. Development of embryos in *Mastacembelus mastacembelus* (Bank & Solender 1794) (Mesopotamian spiny eel) (Mastacembelidae). *Aquac Res*, **37**: 1611-1616.

Sahoo SK, Giri SS, Shaha A, Chandra S, Sahu AK and Sarangi N. 2007. Embryonic development of the spiny eel, *Mastacembelus aculeatus* (Bloch, 1786). *Indian J Fish*, **54(3)**: 333-337.

Serajuddin M and Mustafa S, 1994. Feeding specialization in adult spiny eel, *Mastacembelus armatus*. *Asian Fish Sci*, **7(1)**:63-65.

Suresh VR, Biswas BK, Vinci GK, Mitra K and Mukherjee A. 2006. Biology and fishery of barred spiny eel, *Macrognathus pancalus* (Hamilton). *Acta Ichthyol Piscat*, **36 (1)**: 31-37.

Talwar PK and Jhingran AG. 1991. Inland Fishes of India and Adjacent Countries. Calcutta: Oxford and IBH Publishing.

Umezawa A, Otake T, Hirokawa J, Tsukamoto K and Okiyama M. 1991. Development of the eggs and larvae of the pike eel, *Muraenesox cinereus*. *Jpn J Ichthyol*, **38(1)**:35-40.

Yamamoto K, Yamauchi K and Kasuga S. 1975. On the development of the Japanese eel, *Anguilla japonica*. *Bull Japan Soc Sci Fish*, **41(1)**: 21-28.